arXiv:0911.1100v2 [math.RT] 8 Feb 2011

UNIVERSAL DEFORMATION RINGS OF MODULES OVER FROBENIUS ALGEBRAS

FRAUKE M. BLEHER AND JOSÉ A. VÉLEZ-MARULANDA

Abstract. Let k be a field, and let Λ be a finite dimensional k-algebra. We prove that if Λ is a self-injective algebra, then every finitely generated Λ-module V whose stable endomorphism ring is isomorphic to k has a universal deformation ring $R(\Lambda, V)$ which is a complete local commutative Noetherian k-algebra with residue field k. If Λ is also a Frobenius algebra, we show that $R(\Lambda, V)$ is stable under taking syzygies. We investigate a particular Frobenius algebra Λ_0 of dihedral type, as introduced by Erdmann, and we determine $R(\Lambda_0, V)$ for every finitely generated Λ_0-module V whose stable endomorphism ring is isomorphic to k.

1. Introduction

Let k be a field of arbitrary characteristic, and let W be a complete local commutative Noetherian ring with residue field k. Suppose G is a profinite group and V is a finite dimensional k-vector space with a continuous G-action. If the G-endomorphism ring of V is isomorphic to k and $\mathrm{H}^1(G, \mathrm{End}_k(V))$ is finite dimensional over k, then V has a universal deformation ring $R_W(G, V)$ (see e.g. [11, 17]). The ring $R_W(G, V)$ is a complete local commutative Noetherian W-algebra with residue field k which is universal with respect to isomorphism classes of lifts (i.e. deformations) of V over such W-algebras. Universal deformation rings have become an important tool in number theory, in particular if G is a profinite Galois group (see e.g. [6, 9] and their references). It was shown in [11] that the universal deformation ring $R_W(G, V)$ is isomorphic to the inverse limit of the universal deformation rings $R_W(G_i, V)$ where the G_i range over the (discrete) finite quotients of G through which the action of G on V factors. Thus an important case to consider with respect to deformation rings is when G itself is a finite group, i.e. when kG is a finite dimensional k-algebra.

In this paper, we consider the case when $W = k$ and kG is replaced by a more general finite dimensional k-algebra Λ. Deformations of modules for finite dimensional algebras have been studied by many authors in different contexts (see e.g. [13, 16, 20] and their references). We focus on deformations of modules for such algebras Λ that have properties close to those of group algebras of finite groups. Moreover, our deformations are over arbitrary complete local commutative Noetherian k-algebras with residue field k (see §2 for precise definitions).

Our main motivation to examine this case is as follows. Suppose G is a finite group and V is a finitely generated kG-module. If the endomorphism ring of V is isomorphic to k, or more generally if V is indecomposable, then V belongs to a unique block B of kG. Suppose Λ is a finite dimensional k-algebra that is Morita equivalent to B and V_Λ is the Λ-module that corresponds to V under this Morita equivalence. For example, Λ can be taken to be the basic algebra of B. Then the Morita equivalence provides a correspondence between the deformations of V and the deformations of V_Λ over all complete local commutative Noetherian k-algebras with residue field k, and thus between the deformation rings $R_k(G, V)$ and $R(\Lambda, V_\Lambda)$ (for a precise statement, see Proposition 2.5). Hence this enables us to use methods from the representation theory of finite dimensional algebras to study deformation rings of group representations. This approach has recently led to the solution of various open problems. For example, this was successfully used in [2, 4, 5] to construct

---

2000 *Mathematics Subject Classification.* Primary 16G10; Secondary 16G20, 20C20.

Key words and phrases. Universal deformation rings, Frobenius algebras, stable endomorphism rings.

The first author was supported in part by NSF Grant DMS06-51332. The second author was supported by the Reassigned Time for Research Scholarship of the Office of Academic Affairs at Valdosta State University.

representations whose universal deformation ring is not a complete intersection, thus answering a question posed by Flach [8] in characteristic 2.

Let now Λ be an arbitrary finite dimensional k-algebra and let V be a finitely generated Λ-module. Using Schlessinger's criteria [18], it follows that V always has a versal deformation ring $R(\Lambda, V)$ which is a complete local commutative Noetherian k-algebra with residue field k and that $R(\Lambda, V)$ is universal if the endomorphism ring of V is isomorphic to k. Under the assumption that Λ is a Frobenius algebra, we obtain the following stronger result, which is proved in §2. For a more precise statement, see Theorem 2.6.

Theorem 1.1. *Let Λ be a Frobenius k-algebra and let V be a finitely generated Λ-module whose stable endomorphism ring is isomorphic to k. Then the versal deformation ring $R(\Lambda, V)$ is universal. Moreover, $R(\Lambda, V)$ is isomorphic to $R(\Lambda, V \oplus P)$ for every finitely generated projective Λ-module P, and $R(\Lambda, V)$ is isomorphic to $R(\Lambda, \Omega(V))$ where $\Omega(V)$ is the first syzygy of V.*

Since group algebras of finite groups over k are in particular Frobenius algebras, this generalizes [3, Prop. 2.1 and Cor. 2.5] which give the corresponding results for group algebras. The proof of Theorem 1.1 follows the basic outline of the proofs in [3], but in several instances arguments that are specific to group algebras have to be replaced by arguments that work for arbitrary Frobenius algebras.

In §3, we assume that k is algebraically closed and we consider the particular Frobenius algebra $\Lambda_0 = kQ/I$ where the quiver Q and the ideal I of the path algebra kQ are as in Figure 1. The algebra Λ_0 belongs to the class of algebras of dihedral type which were introduced by Erdmann in [12] to classify all tame blocks of group algebras of finite groups with dihedral defect groups up to Morita equivalence. However, the algebra Λ_0 is not Morita equivalent to a block of a group algebra. Since Λ_0 is a special biserial algebra, all the non-projective indecomposable Λ_0-modules can be described combinatorially as so-called string and band modules (see [7] and §5). The components of the stable Auslander-Reiten quiver $\Gamma_s(\Lambda_0)$ consisting of string modules are two 3-tubes and infinitely many components of type $\mathbb{Z}A_\infty^\infty$, whereas the components consisting of band modules are infinitely many 1-tubes. If M and N are two indecomposable Λ_0-modules belonging to the same component of $\Gamma_s(\Lambda_0)$, we say N is a *successor* of M if there is an irreducible homomorphism $M \to N$.

FIGURE 1. The basic k-algebra $\Lambda_0 = kQ/I$.



$$I = \langle \alpha\lambda, \lambda\xi, \xi\delta, \delta\rho, \rho\beta, \beta\alpha, \alpha^2 - \lambda\delta\beta, \rho^2 - \beta\lambda\delta, \xi^2 - \delta\beta\lambda \rangle.$$

A summary of our main results concerning $\Lambda_0 = kQ/I$ is as follows. The precise statements can be found in Theorem 3.8, Propositions 3.9, 3.10, 3.11, Theorem 3.16 and Proposition 3.17.

Theorem 1.2. *Let $\Lambda_0 = kQ/I$ be as in Figure 1, and suppose $\mathfrak{C}$ is a component of the stable Auslander-Reiten quiver $\Gamma_s(\Lambda_0)$.*

(i) *If $\mathfrak{C}$ is one of the two 3-tubes, then $\Omega(\mathfrak{C})$ is the other 3-tube. There are exactly three Ω^2-orbits of modules in $\mathfrak{C}$ whose stable endomorphism ring is isomorphic to k. If U_0 is a module that belongs to the boundary of $\mathfrak{C}$, then these three Ω^2-orbits are represented by U_0, by a* *successor U_1 of U_0, and by a successor U_2 of U_1 that does not lie in the Ω^2-orbit of U_0. The universal deformation rings are*

$$R(\Lambda_0, U_0) \cong R(\Lambda_0, U_1) \cong k, \qquad R(\Lambda_0, U_2) \cong k[[t]].$$

(ii) *There are infinitely many components of $\Gamma_s(\Lambda_0)$ of type $\mathbb{Z}A_\infty^\infty$ that each contain a module whose stable endomorphism ring is isomorphic to k. If $\mathfrak{C}$ is such a component, then $\mathfrak{C} = \Omega(\mathfrak{C})$ and there are exactly six Ω^2-orbits (resp. exactly three Ω-orbits) of modules in $\mathfrak{C}$ whose stable endomorphism ring is isomorphic to k. If V_0 is a module in $\mathfrak{C}$ of minimal length, then these three Ω-orbits are represented by V_0, by a successor V_1 of V_0 that does not lie in the Ω-orbit of V_0, and by a successor V_2 of V_1 that does not lie in the Ω^2-orbit of V_0. The universal deformation rings are*

$$R(\Lambda_0, V_0) \cong k[[t]]/(t^2), \qquad R(\Lambda_0, V_1) \cong k, \qquad R(\Lambda_0, V_2) \cong k[[t]].$$

(iii) *There are infinitely many 1-tubes of $\Gamma_s(\Lambda_0)$ that each contain a module whose stable endomorphism ring is isomorphic to k. If $\mathfrak{C}$ is such a component, then there is exactly one Ω^2-orbit of modules in $\mathfrak{C}$ whose stable endomorphism ring is isomorphic to k, represented by a module W_0 belonging to the boundary of $\mathfrak{C}$. The universal deformation ring of W_0 is*

$$R(\Lambda_0, W_0) \cong k[[t]].$$

Note that if M is a finitely generated Λ_0-module whose stable endomorphism ring is isomorphic to k, then $M \cong U \oplus P$ where U is an indecomposable Λ_0-module whose stable endomorphism ring is isomorphic to k and P is a projective Λ_0-module. Since by Theorem 1.1, $R(\Lambda_0, M) \cong R(\Lambda_0, U)$, it follows that Theorem 1.2 describes the universal deformation ring for every finitely generated Λ_0-module whose stable endomorphism ring is isomorphic to k.

Theorem 1.2 shows significant differences between Λ_0 and the blocks of group algebras with dihedral defect groups and precisely three isomorphism classes of simple modules which were studied in [1, 2]. Namely for these blocks, there are only finitely many components of the stable Auslander-Reiten quiver containing modules whose stable endomorphism rings are isomorphic to k. Moreover, none of the 1-tubes contain such modules.

The proof of Theorem 1.2 uses the combinatorial description of the indecomposable Λ_0-modules and of the components of the stable Auslander-Reiten quiver of Λ_0. To characterize the infinitely many components in parts (ii) and (iii) of Theorem 1.2, we introduce special words that start and end at the vertex 0 of the quiver Q and which we call words of type 0 (see Definition 3.2). Using an inductive process, we provide a precise description of the modules V_0 from part (ii) (see Theorem 3.8 and Proposition 4.3) and of the modules W_0 from part (iii) (see Theorem 3.16 and Proposition 4.4).

Part of this paper constitutes the Ph.D. thesis of the second author under the supervision of the first author [19].

2. Versal and universal deformation rings

Let k be a field of arbitrary characteristic. Let $\hat{\mathcal{C}}$ be the category of all complete local commutative Noetherian k-algebras with residue field k. The morphisms in $\hat{\mathcal{C}}$ are continuous k-algebra homomorphisms which induce the identity map on k. Let $\mathcal{C}$ be the full subcategory of $\hat{\mathcal{C}}$ of Artinian objects.

Suppose Λ is a finite dimensional k-algebra and V is a finitely generated Λ-module. A *lift* of V over an object R in $\hat{\mathcal{C}}$ is a finitely generated $R \otimes_k \Lambda$-module M which is free over R together with a Λ-module isomorphism $\phi : k \otimes_R M \to V$. Two lifts (M, ϕ) and (M', ϕ') of V over R are *isomorphic* if there exists an $R \otimes_k \Lambda$-module isomorphism $f : M \to M'$ such that $\phi' \circ (k \otimes_R f) = \phi$. The isomorphism class of a lift (M, ϕ) of V over R is denoted by $[M, \phi]$ and called a *deformation* of V over R. We denote the set of all such deformations over R by $\mathrm{Def}_\Lambda(V, R)$. The *deformation functor* $\hat{F}_V : \hat{\mathcal{C}} \to \mathrm{Sets}$ is defined to be the following covariant functor. Let R be an object in $\hat{\mathcal{C}}$ and $\alpha : R \to R'$ a morphism in $\hat{\mathcal{C}}$. Then $\hat{F}_V(R) = \mathrm{Def}_\Lambda(V, R)$ and $\hat{F}_V(\alpha) : \mathrm{Def}_\Lambda(V, R) \to \mathrm{Def}_\Lambda(V, R')$ is defined by $\hat{F}_V(\alpha)([M,\phi]) = [R'\otimes_{R,\alpha} M, \phi_\alpha]$ where $\phi_\alpha : k\otimes_{R'}(R'\otimes_{R,\alpha} M)\to V$ is the composition $k\otimes_{R'}(R'\otimes_{R,\alpha} M)\cong k\otimes_R M \xrightarrow{\phi} V$ of Λ-modules. Let $F_V:\mathcal{C}\to$ Sets be the restriction of $\hat{F}_V$ to the subcategory $\mathcal{C}$ of Artinian objects. Let $k[\epsilon]$, where $\epsilon^2=0$, denote the ring of dual numbers over k. The tangent space of $\hat{F}_V$ and of F_V is defined to be the set $t_V = F_V(k[\epsilon])$. We say that a functor $D:\hat{\mathcal{C}}\to$ Sets is continuous, if for all objects R in $\hat{\mathcal{C}}$ whose maximal ideal is m_R we have $D(R)=\varprojlim_i D(R/m_R^i)$.

Using Schlessinger's criteria [18, Thm. 2.11], it is straightforward to prove the following result:

Proposition 2.1. *The functor F_V has a pro-representable hull $R(\Lambda,V)\in \mathrm{Ob}(\hat{\mathcal{C}})$, as defined in* [18, Def. 2.7]*, and the functor $\hat{F}_V$ is continuous. Moreover, there is a k-vector space isomorphism $t_V\cong \mathrm{Ext}^1_\Lambda(V,V)$. If $\mathrm{End}_\Lambda(V)=k$, then $\hat{F}_V$ is represented by $R(\Lambda,V)$.*

Remark 2.2.
- (i) The first sentence of Proposition 2.1 means that there exists a deformation $[U(\Lambda,V),\phi_U]$ of V over $R(\Lambda,V)$ with the following property. For each $R\in\mathrm{Ob}(\hat{\mathcal{C}})$, the map $\upsilon_R:\mathrm{Hom}_{\hat{\mathcal{C}}}(R(\Lambda,V),R)\to\hat{F}_V(R)$ given by $\alpha\mapsto\hat{F}_V(\alpha)([U(\Lambda,V),\phi_U])$ is surjective, and this map is bijective if R is the ring of dual numbers $k[\epsilon]$.

  If $\hat{F}_V$ is represented by $R(\Lambda,V)$, in the sense that $\hat{F}_V$ is naturally isomorphic to the functor $\mathrm{Hom}_\Lambda(R(\Lambda,V),-)$, then the above map υ_R is bijective for all $R\in\mathrm{Ob}(\hat{\mathcal{C}})$.
- (ii) The isomorphism $t_V\cong\mathrm{Ext}^1_\Lambda(V,V)$ is established in the same manner as in [17, §22]. Namely, given a lift (M,ϕ) of V over $k[\epsilon]$, we have Λ-module isomorphisms $\epsilon\cdot M\cong V$ and $M/\epsilon\cdot M\cong V$. Thus we obtain a short exact sequence of Λ-modules $\mathcal{E}_M: 0\to V\to M\to V\to 0$. The map $s:t_V\to\mathrm{Ext}^1_\Lambda(V,V)$ which sends $[M,\phi]$ to the class $[\mathcal{E}_M]$ in $\mathrm{Ext}^1_\Lambda(V,V)$ is well-defined. It is straightforward to see that s is a k-vector space homomorphism. On the other hand, given an extension $\mathcal{E}: 0\to V\xrightarrow{u_1} V_1\xrightarrow{u_2} V\to 0$ of Λ-modules, we can define a $k[\epsilon]$-module structure on V_1 by letting ϵ act as $u_1\circ u_2$. Thus there is a Λ-module isomorphism $\psi: V_1/\epsilon V_1\to V$, and (V_1,ψ) is a lift of V over $k[\epsilon]$. Sending $[\mathcal{E}]$ to $[V_1,\psi]$ defines the inverse of s.

Definition 2.3. Using the notation from Proposition 2.1 and Remark 2.2(i), the ring $R(\Lambda,V)$ is called the *versal deformation ring* of V and $[U(\Lambda,V),\phi_U]$ is called the *versal deformation* of V. In general, the isomorphism type of $R(\Lambda,V)$ is unique up to a (non-canonical) isomorphism.

If $R(\Lambda,V)$ represents $\hat{F}_V$, then $R(\Lambda,V)$ is called the *universal deformation ring* of V and $[U(\Lambda,V),\phi_U]$ is called the *universal deformation* of V. In this case, $R(\Lambda,V)$ is unique up to a canonical isomorphism.

Remark 2.4. Note that some authors consider a weaker notion of deformations. Namely, let Λ and V be as above, let R be an object in $\hat{\mathcal{C}}$ and let (M,ϕ) be a lift of V over R. Then the isomorphism class $[M]$ of M as an $R\otimes_k\Lambda$-module is called a *weak deformation* of V over R (see e.g. [14, §5.2]). We can also define the *weak deformation functor* $\hat{F}_V^w:\hat{\mathcal{C}}\to$ Sets which sends an object R in $\hat{\mathcal{C}}$ to the set of weak deformations of V over R and a morphism $\alpha:R\to R'$ in $\hat{\mathcal{C}}$ to the map $\hat{F}_V^w(\alpha):\hat{F}_V^w(R)\to\hat{F}_V^w(R')$ which is defined by $\hat{F}_V^w(\alpha)([M])=[R'\otimes_{R,\alpha}M]$.

In general, a weak deformation of V over R identifies more lifts than a deformation of V over R that respects the isomorphism ϕ of a representative (M,ϕ). However, if Λ is self-injective and the stable endomorphism ring $\underline{\mathrm{End}}_\Lambda(V)$ is isomorphic to k, then the two deformation functors $\hat{F}_V$ and $\hat{F}_V^w$ are naturally isomorphic (see Theorem 2.6(i)).

We now prove that Morita equivalences preserve versal deformation rings. Recall that two finitely generated k-algebras Λ and Λ' are said to be Morita equivalent if they have equivalent module categories. By the Morita theorems (see for example [10, §3D]), Λ and Λ' are Morita equivalent if and only if there exist a Λ'-Λ-bimodule P and a Λ-Λ'-bimodule Q such that P and Q are finitely generated projective both as left and as right modules and $Q\otimes_{\Lambda'}P\cong\Lambda$ as Λ-Λ-bimodules and $P\otimes_\Lambda Q\cong\Lambda'$ as Λ'-Λ'-bimodules. In this situation, we say that P and Q induce a Morita equivalence between Λ and Λ'. Note that if P and Q are such bimodules, then $P\otimes_\Lambda -$ and $Q\otimes_{\Lambda'} -$ give mutually inverse equivalences between the module categories of Λ and Λ'.

Proposition 2.5. *Let Λ and Λ' be finitely generated k-algebras. Suppose that P is a Λ'-Λ-bimodule and Q is a Λ-Λ'-bimodule which induce a Morita equivalence between Λ and Λ'. Let V be a finitely generated Λ-module, and define $V' = P \otimes_\Lambda V$. Then the versal deformation rings $R(\Lambda, V)$ and $R(\Lambda', V')$ are isomorphic in $\hat{\mathcal{C}}$.*

Proof. Suppose $R \in \mathrm{Ob}(\mathcal{C})$ is Artinian, and define $R\Lambda = R \otimes_k \Lambda$ and $R\Lambda' = R \otimes_k \Lambda'$. Then $P_R = R \otimes_k P$ is finitely generated projective as a left $R\Lambda'$-module and as a right $R\Lambda$-module, and $Q_R = R \otimes_k Q$ is finitely generated projective as a left $R\Lambda$-module and as a right $R\Lambda'$-module. We have

$$P_R \otimes_{R\Lambda} Q_R \cong R \otimes_k (P \otimes_\Lambda Q) \cong R\Lambda \quad \text{as } R\Lambda\text{-}R\Lambda\text{-bimodules}$$

and

$$Q_R \otimes_{R\Lambda'} P_R \cong R \otimes_k (Q \otimes_{\Lambda'} P) \cong R\Lambda' \quad \text{as } R\Lambda'\text{-}R\Lambda'\text{-bimodules.}$$

In particular, $P_R \otimes_{R\Lambda} -$ and $Q_R \otimes_{R\Lambda'} -$ give mutually inverse equivalences between the module categories of $R\Lambda$ and $R\Lambda'$.

Let now (M, ϕ) be a lift of V over R. Then M is a finitely generated $R\Lambda$-module. Define $M' = P_R \otimes_{R\Lambda} M$. Since P_R is a finitely generated projective right $R\Lambda$-module and since M is a finitely generated free R-module, it follows that M' is a finitely generated projective, and hence free, R-module. Moreover,

$$k \otimes_R M' = k \otimes_R (P_R \otimes_{R\Lambda} M) = P \otimes_\Lambda (k \otimes_R M) \xrightarrow{P\otimes_\Lambda \phi} P \otimes_\Lambda V = V'. \tag{2.1}$$

This means that $(M', \phi') = (P_R \otimes_{R\Lambda} M, P \otimes_\Lambda \phi)$ is a lift of V' over R. We therefore obtain for all $R \in \mathrm{Ob}(\mathcal{C})$ a well-defined map

$$\tau_R : \mathrm{Def}_\Lambda(V, R) \to \mathrm{Def}_{\Lambda'}(V', R).$$

Since $P_R \otimes_{R\Lambda} -$ and $Q_R \otimes_{R\Lambda'} -$ give mutually inverse equivalences between the module categories of $R\Lambda$ and $R\Lambda'$, it follows that τ_R is bijective. It is straightforward to check that τ_R is natural with respect to homomorphisms $\alpha : R \to R'$ in $\mathcal{C}$. Since the deformation functors $\hat{F}_V$ and $\hat{F}_{V'}$ are continuous, this implies that they are naturally isomorphic. Hence the versal deformation rings $R(\Lambda, V)$ and $R(\Lambda', V')$ are isomorphic in $\hat{\mathcal{C}}$. □

In [3], it was proved that if Λ is the group algebra kG of a finite group G, then a finitely generated Λ-module V has a universal deformation ring if the stable endomorphism ring $\underline{\mathrm{End}}_\Lambda(V)$ is isomorphic to k. Moreover, in this case $R(\Lambda, V) \cong R(\Lambda, \Omega(V))$, where $\Omega(V)$ is the first syzygy of V, i.e. $\Omega(V)$ is the kernel of a projective cover $P_V \to V$. We want to generalize this result to arbitrary Frobenius algebras. Recall that a finite dimensional k-algebra Λ is a Frobenius algebra if and only if there exists a non-degenerate associative bilinear form $\theta : \Lambda \times \Lambda \to k$. This is equivalent to the statement that the right Λ-modules Λ_Λ and $({}_\Lambda\Lambda)^* = \mathrm{Hom}_k({}_\Lambda\Lambda, k)$ are isomorphic. In particular, the group algebra kG is a Frobenius algebra. By [10, Prop. 9.9], every Frobenius algebra is self-injective.

Theorem 2.6. *Let Λ be a finite dimensional self-injective k-algebra, and suppose V is a finitely generated Λ-module whose stable endomorphism ring $\underline{\mathrm{End}}_\Lambda(V)$ is isomorphic to k.*

- (i) *The deformation functor $\hat{F}_V$ is naturally isomorphic to the weak deformation functor $\hat{F}_V^w$ from Remark 2.4.*
- (ii) *The module V has a universal deformation ring $R(\Lambda, V)$.*
- (iii) *If P is a finitely generated projective Λ-module, then $\underline{\mathrm{End}}_\Lambda(V \oplus P) \cong k$ and $R(\Lambda, V) \cong R(\Lambda, V \oplus P)$.*
- (iv) *If Λ is also a Frobenius algebra, then $\underline{\mathrm{End}}_\Lambda(\Omega(V)) \cong k$ and $R(\Lambda, V) \cong R(\Lambda, \Omega(V))$.*

Proof. Suppose $\underline{\mathrm{End}}_\Lambda(V) \cong k$. Then $\underline{\mathrm{End}}_\Lambda(V \oplus P) \cong k$ for every finitely generated projective Λ-module P. Since Λ is self-injective, Ω induces a self-equivalence of the stable module category Λ-$\underline{\mathrm{mod}}$. Hence also $\underline{\mathrm{End}}_\Lambda(\Omega(V)) \cong k$.

We want to follow the basic outline of the proofs of [3, Lemma 2.3 and Props. 2.4 and 2.6]. Because the functor $\hat{F}_V$ (resp. $\hat{F}_V^w$) is continuous, most of the arguments can be carried out for the restriction F_V (resp. F_V^w) of $\hat{F}_V$ (resp. $\hat{F}_V^w$) to the subcategory $\mathcal{C}$ of $\hat{\mathcal{C}}$ of Artinian objects. If $R \in \mathrm{Ob}(\mathcal{C})$ is Artinian, we define $R\Lambda = R \otimes_k \Lambda$.

Let $\pi : R \to R_0$ be a surjection in $\mathcal{C}$, and let Q_0 be a finitely generated projective $R_0\Lambda$-module. Since $R\Lambda$ is Artinian, Q_0 has a projective $R\Lambda$-module cover $\mathrm{Proj}_R(Q_0)$, which is unique up to isomorphism. In particular, $R_0 \otimes_{R,\pi} \mathrm{Proj}_R(Q_0) \cong Q_0$ as $R_0\Lambda$-modules. We first show in Claims 1 and 2 below that certain homomorphisms between $R_0\Lambda$-modules which factor through projective modules can be lifted to $R\Lambda$-module homomorphisms. Since R, R_0 are Artinian, we can use an inductive argument, assuming that $\pi : R \to R_0$ is a small extension, i.e. the kernel of π is a principal ideal tR annihilated by the maximal ideal m_R of R.

Claim 1. Let $\pi : R \to R_0$ be a surjection in $\mathcal{C}$. Let M, Q (resp. M_0, Q_0) be finitely generated $R\Lambda$-modules (resp. $R_0\Lambda$-modules) and assume that Q (resp. Q_0) is projective. Suppose there are $R_0\Lambda$-module isomorphisms $g : R_0 \otimes_{R,\pi} M \to M_0$, $h : R_0 \otimes_{R,\pi} Q \to Q_0$. If $\nu_0 \in \mathrm{Hom}_{R_0\Lambda}(M_0, Q_0)$, then there exists $\nu \in \mathrm{Hom}_{R\Lambda}(M, Q)$ with $\nu_0 = h \circ (R_0 \otimes_{R,\pi} \nu) \circ g^{-1}$.

Proof of Claim 1. As noted in the paragraph before the statement of Claim 1, we can assume that $\pi : R \to R_0$ is a small extension whose kernel is tR. Tensoring $0 \to tR \to R \xrightarrow{\pi} R_0 \to 0$ with Q over R, we obtain a short exact sequence of $R\Lambda$-modules

$$0 \to tQ \to Q \xrightarrow{h\circ\tau_Q} Q_0 \to 0 \tag{2.2}$$

where $\tau_Q : Q \to R_0 \otimes_{R,\pi} Q$ is the natural surjection. Applying $\mathrm{Hom}_{R\Lambda}(M, -)$ to (2.2), we obtain a long exact sequence

$$\cdots \to \mathrm{Hom}_{R\Lambda}(M, Q) \xrightarrow{(h\circ\tau_Q)_*} \mathrm{Hom}_{R\Lambda}(M, Q_0) \to \mathrm{Ext}^1_{R\Lambda}(M, tQ) \to \cdots \tag{2.3}$$

Because $tQ \cong k \otimes_R Q$ is a finitely generated projective Λ-module, and hence an injective Λ-module, it follows that $\mathrm{Ext}^1_{R\Lambda}(M, tQ) \cong \mathrm{Ext}^1_{\Lambda}(k \otimes_R M, tQ) = 0$. Thus the map $(h \circ \tau_Q)_*$ in (2.3) is surjective. Since the natural surjection $\tau_M : M \to R_0 \otimes_{R,\pi} M$ induces an isomorphism $\mathrm{Hom}_{R_0\Lambda}(M_0, Q_0) \xrightarrow{(g\circ\tau_M)^*} \mathrm{Hom}_{R\Lambda}(M, Q_0)$, this implies that there exists an $R\Lambda$-module homomorphism $\nu : M \to Q$ such that $\nu_0 = h \circ (R_0 \otimes_{R,\pi} \nu) \circ g^{-1}$. This proves Claim 1.

As noted in the paragraph before Claim 1, finitely generated projective $R_0\Lambda$-modules can be lifted to finitely generated projective $R\Lambda$-modules. We can use this together with Claim 1 to prove the following result.

Claim 2. Let $\pi : R \to R_0$ be a surjection in $\mathcal{C}$. Let M (resp. M_0) be a finitely generated $R\Lambda$-module (resp. $R_0\Lambda$-module) such that there is an $R_0\Lambda$-module isomorphism $g : R_0 \otimes_{R,\pi} M \to M_0$. Suppose $\sigma_0 \in \mathrm{End}_\Lambda(M_0)$ factors through a projective $R_0\Lambda$-module. Then there exists $\sigma \in \mathrm{End}_{R\Lambda}(M)$ such that σ factors through a projective $R\Lambda$-module and $\sigma_0 = g \circ (R_0 \otimes_{R,\pi} \sigma) \circ g^{-1}$.

We next show in Claim 3 that if (M, ϕ) is a lift of V over an object R in $\mathcal{C}$, then the deformation $[M, \phi]$ does not depend on the particular choice of the Λ-module isomorphism. This implies part (i) of Theorem 2.6 since $\hat{F}_V$ and $\hat{F}_V^w$ are continuous. In other words, the deformation functor $\hat{F}_V$ can be identified with the deformation functor considered in [3].

Claim 3. Let R be an Artinian object in $\mathcal{C}$, and let (M, ϕ) and (M', ϕ') be two lifts of V over R. If there is an $R\Lambda$-module isomorphism $f : M \to M'$, then there exists an $R\Lambda$-module isomorphism $f' : M \to M'$ such that $\phi' \circ (k \otimes_R f') = \phi$. In other words, $[M, \phi] = [M', \phi']$.

Proof of Claim 3. Let $\overline{f}'' = \phi \circ (k \otimes_R f)^{-1} \circ (\phi')^{-1} \in \mathrm{End}_\Lambda(V)$. Since $\underline{\mathrm{End}}_\Lambda(V) = k$ by assumption, there exists a scalar $\overline{s_f} \in k$ and a homomorphism $\overline{\sigma}_f \in \mathrm{End}_\Lambda(V)$ which factors through a projective Λ-module such that $\overline{f}'' = \overline{s_f} \cdot \mathrm{id}_V + \overline{\sigma}_f$. By Claim 2, there exists $\sigma_f \in \mathrm{End}_{R\Lambda}(M)$ which factors through a projective $R\Lambda$-module such that $\overline{\sigma}_f = \phi \circ (k \otimes_R \sigma_f) \circ \phi^{-1}$. Moreover, there exists $s_f \in R$ such that $k \otimes_R s_f = \overline{s_f}$. Let $f'' = s_f \cdot \mathrm{id}_M + \sigma_f$. Then $k \otimes_R f'' = \phi^{-1} \circ \overline{f}'' \circ \phi$ is an automorphism of

$k\otimes_R M$, which implies by Nakayama's Lemma that f'' is an automorphism of M. Define $f' = f\circ f''$. Then $\phi'\circ(k\otimes_R f') = \phi$, which proves Claim 3.

We can now use Claims 2 and 3 together with analogous arguments to the ones used in the proof of [3, Lemma 2.3] to prove the following result.

Claim 4. If (M,ϕ) is a lift of V over an Artinian object R in $\mathcal{C}$, then the ring homomorphism $\sigma_M : R \to \underline{\mathrm{End}}_{R\Lambda}(M)$ coming from the action of R on M via scalar multiplication is surjective.

We next prove part (ii) of Theorem 2.6. Because of Proposition 2.1, it suffices to verify Schlessinger's criterion (H_4) in [18, Thm. 2.11], which is a consequence of the following result.

Claim 5. Let $\pi : R' \to R$ be a small extension of Artinian objects in $\mathcal{C}$. Then the natural map $F_V(R'\times_R R') \to F_V(R')\times_{F_V(R)} F_V(R')$ is injective.

Proof of Claim 5. Let $\alpha_1 : R'\times_R R' \to R'$ (resp. $\alpha_2 : R'\times_R R' \to R'$) be the natural surjection onto the first (resp. second) component of $R'\times_R R'$ such that $\pi\circ\alpha_1 = \pi\circ\alpha_2$. Let (M_1,ϕ_1) and (M_2,ϕ_2) be lifts of V over $R'\times_R R'$. Suppose there are $R'\Lambda$-module isomorphisms $f_i : R'\otimes_{R'\times_R R',\alpha_i} M_1 \to R'\otimes_{R'\times_R R',\alpha_i} M_2$ for $i=1,2$. Then $g_R = (R\otimes_{R',\pi} f_2)^{-1}\circ(R\otimes_{R',\pi} f_1)$ is an $R\Lambda$-module automorphism of $R\otimes_{R'\times_R R',\pi\circ\alpha_2} M_1$. By Claims 2 and 4, it follows that there exists an $R'\Lambda$-module automorphism g of $R'\otimes_{R'\times_R R',\alpha_2} M_1$ with $R\otimes_{R',\pi} g = g_R$. Replacing f_2 by $f_2\circ g$, we have $R\otimes_{R',\pi} f_1 = R\otimes_{R',\pi} f_2$. Therefore, the pair (f_1,f_2) defines an $(R'\times_R R')\Lambda$-module isomorphism $f : M_1 \to M_2$. By Claim 3, this implies that $[M_1,\phi_1] = [M_2,\phi_2]$, which proves Claim 5.

For part (iii) of Theorem 2.6, we need the following result.

Claim 6. Suppose P is a finitely generated projective Λ-module and there is a commutative diagram of finitely generated $R\Lambda$-modules

$$\begin{tikzcd} 0 \arrow[r] & \mathrm{Proj}_R(P) \arrow[r, "\alpha"] \arrow[d] & T \arrow[r, "\beta"] \arrow[d] & C \arrow[r] \arrow[d] & 0 \\ 0 \arrow[r] & P \arrow[r, "\overline{\alpha}"] & k\otimes_R T \arrow[r, "\overline{\beta}"] & k\otimes_R C \arrow[r] & 0 \end{tikzcd} \tag{2.4}$$

in which T and C are free over R and the bottom row arises by tensoring the top row with k over R and identifying P with $k\otimes_R \mathrm{Proj}_R(P)$. Then the top row of (2.4) splits as a sequence of $R\Lambda$-modules.

Proof of Claim 6. Since Λ is self-injective, P is an injective Λ-module. Thus there exists a Λ-module homomorphism $\overline{\omega} : k\otimes_R T \to P$ with $\overline{\omega}\circ\overline{\alpha} = \mathrm{id}_P$. By Claim 1, there exists an $R\Lambda$-module homomorphism $\omega : T \to \mathrm{Proj}_R(P)$ such that $k\otimes_R\omega = \overline{\omega}$. Hence $k\otimes_R(\omega\circ\alpha) = \overline{\omega}\circ\overline{\alpha} = \mathrm{id}_P$. Using Nakayama's Lemma, this implies that $\omega\circ\alpha$ is an $R\Lambda$-module automorphism of $\mathrm{Proj}_R(P)$. This proves Claim 6.

Part (iii) of Theorem 2.6 follows now by using Claim 6 together with analogous arguments to the ones used in the proofs of [3, Prop. 2.6 and Cor. 2.7].

For the proof of part (iv) of Theorem 2.6, we assume that Λ is a Frobenius algebra. We need the following result.

Claim 7. Let R be an Artinian object in $\mathcal{C}$, and let Q be a finitely generated projective left $R\Lambda$-module. Then $Q^* = \mathrm{Hom}_{R\Lambda}(Q,R)$ is a projective right $R\Lambda$-module.

Proof of Claim 7. Consider the additive map

$$\begin{array}{rcl} R\otimes_k \mathrm{Hom}_k({}_\Lambda\Lambda, k) & \to & \mathrm{Hom}_R({}_{R\otimes_k\Lambda}(R\otimes_k\Lambda), R) = \mathrm{Hom}_R({}_{R\Lambda}(R\Lambda), R) \\ b\otimes f & \mapsto & (a\otimes\lambda \mapsto abf(\lambda)). \end{array} \tag{2.5}$$

By [10, Thm. 2.38], (2.5) is an R-module isomorphism. It is straightforward to check that (2.5) is in fact an isomorphism of right $R\Lambda$-modules. Since Λ is a Frobenius algebra, there is an isomorphism

$\Lambda_\Lambda \cong ({}_\Lambda\Lambda)^* = \mathrm{Hom}_k({}_\Lambda\Lambda, k)$ of right Λ-modules. Therefore,

$$({}_{R\Lambda}(R\Lambda))^* = \mathrm{Hom}_R({}_{R\Lambda}(R\Lambda), R) \cong R \otimes_k ({}_\Lambda\Lambda)^* \cong R \otimes_k (\Lambda_\Lambda) \cong (R\Lambda)_{R\Lambda}$$

as right $R\Lambda$-modules. By assumption, Q is a finitely generated projective left $R\Lambda$-module. Hence there exists a left $R\Lambda$-module T and an integer n such that $Q \oplus T \cong ({}_{R\Lambda}(R\Lambda))^n$. Since Q and T are free over R, they are $R\Lambda$-lattices. Thus we have

$$Q^* \oplus T^* \cong (Q \oplus T)^* \cong \left(({}_{R\Lambda}(R\Lambda))^*\right)^n \cong ((R\Lambda)_{R\Lambda})^n .$$

Hence Q^* is a projective right $R\Lambda$-module, which proves Claim 7.

Consider the short exact sequence $0 \to \Omega(V) \xrightarrow{\overline{\varphi}} P_V \to V \to 0$ coming from the definition of $\Omega(V)$. Let (U, ψ) be a lift of $\Omega(V)$ over an Artinian object R in $\mathcal{C}$. Since $\mathrm{Proj}_R(P_V)$ is a projective $R\Lambda$-module with $k \otimes_R \mathrm{Proj}_R(P_V) \cong P_V$, we can use Claim 1 to lift $\overline{\varphi}$ to an $R\Lambda$-module homomorphism $\varphi : U \to \mathrm{Proj}_R(P_V)$. Using this together with Claim 7, we can argue analogously to the proofs of [3, Prop. 2.4 and Cor. 2.5] to prove part (iv) of Theorem 2.6.

This completes the proof of Theorem 2.6. □

3. A PARTICULAR ALGEBRA OF DIHEDRAL TYPE

For the remainder of this article, let k be an algebraically closed field of arbitrary characteristic and let Λ_0 be the basic k-algebra $\Lambda_0 = kQ/I$ where Q and I are as in Figure 1. The algebra Λ_0 is one of the algebras of dihedral type studied by Erdmann in [12]. In particular, Λ_0 is symmetric, and hence a Frobenius algebra. By [12, Lemma IX.5.4], Λ_0 is not Morita equivalent to a block of a group algebra. We denote the irreducible Λ_0-modules by S_0, S_1 and S_2, or, using shorthand, by 0, 1 and 2. The radical series of the projective indecomposable Λ_0-modules can be described by the following pictures:

$$P_0 = \begin{array}{ccc} & 0 & \\ 0 & & \begin{array}{c}1\\2\end{array} \\ & 0 & \end{array}, \qquad P_1 = \begin{array}{ccc} & 1 & \\ 1 & & \begin{array}{c}2\\0\end{array} \\ & 1 & \end{array}, \qquad P_2 = \begin{array}{ccc} & 2 & \\ 2 & & \begin{array}{c}0\\1\end{array} \\ & 2 & \end{array}. \tag{3.1}$$

Since Λ_0 is a special biserial algebra, all indecomposable non-projective Λ_0-modules are either string or band modules (see [7]). We give an introduction to the representation theory of Λ_0 in an appendix in §5, where we introduce in particular words, strings, bands and canonical k-bases for string and band modules. Moreover, we describe the components of the stable Auslander-Reiten quiver using hooks and cohooks and give a description of the homomorphisms between string and band modules as determined in [15].

Our goal in this section is to determine all string and band modules V for Λ_0 whose stable endomorphism rings are isomorphic to k and to compute their universal deformation rings $R(\Lambda_0, V)$. Because of the symmetries that both the quiver Q and the relations I satisfy, we will make use of the following permutations of the vertices and arrows of Q.

Definition 3.1. (i) For $u \in \{0, 1, 2\}$, define ν_u to be the following permutation of the vertices and arrows of Q:

$$\nu_0 = (1,2)(\rho,\xi)(\beta,\lambda), \quad \nu_1 = (0,2)(\alpha,\xi)(\beta,\delta), \quad \text{and} \quad \nu_2 = (0,1)(\alpha,\rho)(\lambda,\delta).$$

Note that ν_0, ν_1 and ν_2 are not quiver automorphisms of Q. If $S = w_1 w_2 \cdots w_n$ is a word of length $n \geq 1$ representing a string, let $u_S = e(S)$ and define

$$\nu_{u_S}(S) = \nu_{u_S}(w_1)^{-1} \nu_{u_S}(w_2)^{-1} \cdots \nu_{u_S}(w_n)^{-1}.$$

(ii) Let θ be the quiver automorphism of Q which acts on the vertices and arrows of Q as the permutation

$$\theta = (0,1,2)(\alpha,\rho,\xi)(\beta,\delta,\lambda).$$

If $S = w_1 w_2 \cdots w_n$ is a word of length $n \geq 1$ representing a string, define

$$\theta(S) = \theta(w_1)\theta(w_2) \cdots \theta(w_n).$$

The following words starting and ending at the vertex 0 will play a special role when discussing both string and band modules.

Definition 3.2. (i) Define $\underline{x} = \lambda\xi^{-1}\delta\beta\alpha^{-1}$ and $\underline{y} = \lambda\delta\rho^{-1}\beta\alpha^{-1}$.

(ii) We say a word Z is a *word of type* 0 if $Z = 1_0$ or

$$Z = \underline{z}_0\underline{z}_1 \cdots \underline{z}_{n-1}$$

where $n \geq 1$ and $\underline{z}_0, \ldots, \underline{z}_{n-1} \in \{\underline{x}, \underline{y}\}$. We call n the *type-0-length* of Z, where $n = 0$ corresponds to $Z = 1_0$.

(iii) Let $Z = \underline{z}_0\underline{z}_1 \cdots \underline{z}_{n-1}$ be a word of type 0. A *subpattern* U of Z is a subword of Z such that either $U = 1_0$ or $U = \underline{z}_i\underline{z}_{i+1} \cdots \underline{z}_j$ for some $0 \leq i \leq j \leq n-1$. The *inverse pattern* U^- of U is defined to be $U^- = 1_0$ if $U = 1_0$ and $U^- = \underline{z}_j\underline{z}_{j-1} \cdots \underline{z}_i$ if $U = \underline{z}_i\underline{z}_{i+1} \cdots \underline{z}_j$ for some $0 \leq i \leq j \leq n-1$.

(iv) Let $Z = \underline{z}_0 \cdots \underline{z}_{n-1}$ be a word of type 0 of type-0-length $n \geq 1$. For $0 \leq i \leq n-1$, we define the i^{th} *rotation of Z of type* 0 to be

$$\rho_i^{(0)}(Z) = \underline{z}_i\underline{z}_{i+1} \cdots \underline{z}_{n-1}\underline{z}_0 \cdots \underline{z}_{i-1}.$$

Define $\sim_{r,(0)}$ to be the equivalence relation on all words of type 0 of type-0-length at least 1 such that $Z \sim_{r,(0)} Z'$ if and only if $Z = \rho_i^{(0)}(Z')$ for some i.

3.1. Stable endomorphism rings and universal deformation rings of string modules for Λ_0.

In this section, we describe all string modules $M(S)$ for Λ_0 whose stable endomorphism rings are isomorphic to k and determine their universal deformation rings.

The following result is an easy consequence of [15] (see Remark 5.4) and the symmetric shapes of the radical series of P_0, P_1 and P_2 as given in (3.1).

Lemma 3.3. *Let $S = w_1w_2 \cdots w_n$ be a word of length $n \geq 1$ representing a string and let $\nu_{u_S}(S)$ be as in Definition 3.1(i). Then $\nu_{u_S}(S)$ is a word representing a string. Moreover,*

$$\begin{aligned} \dim_k \underline{\mathrm{End}}_{\Lambda_0}(M(S)) &= \dim_k \underline{\mathrm{End}}_{\Lambda_0}(M(\nu_{u_S}(S))), \quad \textit{and} \\ \dim_k \mathrm{Ext}^1_{\Lambda_0}(M(S), M(S)) &= \dim_k \mathrm{Ext}^1_{\Lambda_0}(M(\nu_{u_S}(S)), M(\nu_{u_S}(S))). \end{aligned}$$

For the description of all the string modules whose stable endomorphism rings are isomorphic to k, we need to introduce special strings, which we call strings of type $\beta\alpha^{-1}\lambda$ and which are defined using the words of type 0 introduced in Definition 3.2.

Definition 3.4. (i) We say a word S is the *standard representative of a string of type $\beta\alpha^{-1}\lambda$* if

$$S = \beta\alpha^{-1}Z\,\lambda$$

where Z is a word of type 0, as defined in Definition 3.2(ii). A subword U of S is called a *subpattern* of S if U is a subpattern of Z, as defined in Definition 3.2(iii). The *inverse pattern* U^- of U is also defined as in Definition 3.2(iii). We define $S^- = \beta\alpha^{-1}Z^-\lambda$.

(ii) Define Ξ to be the permutation $\Xi = (\underline{x}, \underline{y})$. Define $\Xi(1_0) = 1_0$. If $Z = \underline{z}_0 \cdots \underline{z}_{n-1}$ is a word of type 0 of type-0-length $n \geq 1$, define $\Xi(Z) = \Xi(\underline{z}_0) \cdots \Xi(\underline{z}_{n-1})$. If $S = \beta\alpha^{-1}Z\lambda$ is the standard representative of a string of type $\beta\alpha^{-1}\lambda$, define $\Xi(S) = \beta\alpha^{-1}\Xi(Z)\lambda$ and $\Xi(S^-) = \beta\alpha^{-1}\Xi(Z^-)\lambda$.

Remark 3.5. Let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$, and let S^- and Ξ be as in Definition 3.4. Moreover, let ν_0, ν_1, ν_2 and θ be as in Definition 3.1. Then $\theta(\nu_1(S)) = \nu_2(\theta(S)) = \Xi(S^-)^{-1}$, and hence $\Xi(S^-) = \theta^2(\nu_2(S^{-1}))$. Moreover, $\nu_2(S^{-1}) = \left(\theta^2(\nu_1(S))\right)^{-1}$.

Lemma 3.6. *Let $S = \beta\alpha^{-1}\underline{z}_0 \cdots \underline{z}_{n-1}\,\lambda$ be the standard representative of a string of type $\beta\alpha^{-1}\lambda$ as in Definition 3.4. The stable endomorphism ring of $M(S)$, $\underline{\mathrm{End}}_{\Lambda_0}(M(S))$, is isomorphic to k if and only if S has the following property:*

(+) *If there are subpatterns* U, A_1, A_2 *of* S *such that*

$$S = \beta\alpha^{-1}\, U\, \underline{x}\, A_2\, \lambda \quad \text{or} \quad S = \beta\alpha^{-1}\, A_1\, \underline{x}\, U\, \underline{x}\, A_2\, \lambda \quad \text{or} \quad S = \beta\alpha^{-1}\, A_1\, \underline{x}\, U\, \lambda,$$

then for all subpatterns B_1, B_2 *of* S, *we have*

$$S \neq \beta\alpha^{-1}\, U\, \underline{y}\, B_2\, \lambda \quad \text{and} \quad S \neq \beta\alpha^{-1}\, B_1\, \underline{y}\, U\, \underline{y}\, B_2\, \lambda \quad \text{and} \quad S \neq \beta\alpha^{-1}\, B_1\, \underline{y}\, U\, \lambda.$$

Moreover, if the stable endomorphism ring of $M(S)$ *has* k*-dimension at least* 2, *then all modules in the stable Auslander-Reiten component containing* $M(S)$ *have this property.*

Proof. Since $\underline{\mathrm{End}}_{\Lambda_0}(M(\beta\alpha^{-1}\lambda)) \cong k$ and $\beta\alpha^{-1}\lambda$ has property (+), Lemma 3.6 follows for $n = 0$. For the remainder of the proof, let $S = \beta\alpha^{-1}\underline{z}_0 \cdots \underline{z}_{n-1}\,\lambda$ be a word representing a string of type $\beta\alpha^{-1}\lambda$ with $n \geq 1$. We analyze the different possibilities for $M(S)$ such that $\underline{\mathrm{End}}_{\Lambda_0}(M(S))$ has k-dimension at least 2.

Suppose first that there exists a string $T \not\sim_s S$ of length at least 2 such that $M(S)$ has a canonical endomorphism α_T, as defined in Remark 5.4, factoring through $M(T)$. In particular, $\underline{\mathrm{End}}_{\Lambda_0}(M(S))$ has k-dimension at least 2. It follows from the definition of strings of type $\beta\alpha^{-1}\lambda$ that

$$T \sim_s \beta\alpha^{-1}\, U\, \lambda$$

for some subpattern U of S and that there exist subpatterns A_1, A_2, B_1, B_2 of S satisfying both $(*)$ and $(**)$, where

$(*)$ $S = \beta\alpha^{-1}\, U\, \underline{x}\, A_2\, \lambda$ or $S = \beta\alpha^{-1}\, A_1\, \underline{x}\, U\, \underline{x}\, A_2\, \lambda$ or $S = \beta\alpha^{-1}\, A_1\, \underline{x}\, U\, \lambda$,
$(**)$ $S = \beta\alpha^{-1}\, U\, \underline{y}\, B_2\, \lambda$ or $S = \beta\alpha^{-1}\, B_1\, \underline{y}\, U\, \underline{y}\, B_2\, \lambda$ or $S = \beta\alpha^{-1}\, B_1\, \underline{y}\, U\, \lambda$.

On the other hand, if S satisfies $(*)$ and $(**)$, then $M(S)$ has a canonical endomorphism factoring through a string module that is not isomorphic to $M(S)$ with at least three composition factors. It is straightforward to see that then also the modules $M(S_{h\cdots h})$ and $M({}_{h\cdots h}S)$ each have a non-identity canonical endomorphism factoring through a string module with at least three composition factors.

Next let T be a string of length 0 or 1 such that $M(S)$ has a canonical endomorphism α_T factoring through $M(T)$. Then either $T = \delta$ or $T = 1_u$ for some $u \in \{0, 1, 2\}$. Analyzing these cases further, it follows that α_T does not factor through a projective Λ_0-module only if S satisfies $(*)$ and $(**)$ for certain subpatterns U, A_1, A_2, B_1, B_2 of S. This completes the proof of Lemma 3.6. □

Remark 3.7. Let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$, and let S^- and Ξ be as in Definition 3.4. If S satisfies condition (+) from Lemma 3.6, then $S = S^-$. Moreover, S satisfies condition (+) if and only if $\Xi(S)$ satisfies condition (+).

Theorem 3.8. *Let* $\mathfrak{C}$ *be a component of the stable Auslander-Reiten quiver* $\Gamma_s(\Lambda_0)$ *consisting of string modules. Then* $\mathfrak{C}$ *contains a module whose stable endomorphism ring is isomorphic to* k *if and only if either*

(i) $\mathfrak{C}$ *is a* 3*-tube, or*
(ii) $\mathfrak{C}$ *contains a simple* Λ_0*-module, or*
(iii) *there exist* $i \in \{0, 1, 2\}$ *and a string of type* $\beta\alpha^{-1}\lambda$ *whose standard representative* S *satisfies condition* (+) *from Lemma 3.6 such that* $\mathfrak{C}$ *contains* $M(\theta^i(S))$.

Proof. Since one of the 3-tubes contains $M(\alpha)$ and the other 3-tube contains $M(\delta\beta)$, it follows that if $\mathfrak{C}$ is as in (i), (ii) or (iii), then $\mathfrak{C}$ contains a module whose stable endomorphism ring is isomorphic to k.

Conversely, let $\mathfrak{C}$ be a component of string modules containing a module whose stable endomorphism ring is isomorphic to k, and suppose $\mathfrak{C}$ is not a 3-tube. Then $\mathfrak{C}$ is of type $\mathbb{Z}A_\infty^\infty$. Let T be a word representing a string such that $M(T)$ has minimal length in $\mathfrak{C}$.

If T has length 0, 1 or 2, it follows that $M(T)$ lies in the same component as a simple Λ_0-module, and hence $M(T)$ is itself simple.

Now suppose T is a string of length at least 3. Let $u_T = e(T)$, let ν_{u_T} and θ be as in Definition 3.1, and let $i_T \in \{0, 1, 2\}$ be such that $\theta^{i_T}(u_T) = 1$. Then there exists $j \in \{0, 1\}$ such that

$$\tilde{S} \overset{\text{def}}{=} \theta^{i_T}(\nu_{u_T}^{j}(T)) = \beta\alpha^{-1}\lambda\cdots . \tag{3.2}$$

Suppose first that $\tilde{S}$ in (3.2) is not the standard representative of a string of type $\beta\alpha^{-1}\lambda$. Then there exists a word $Z = \underline{z}_0 \cdots \underline{z}_{n-1}$ of type 0 of maximal type-0-length $n \geq 0$ such that $\tilde{S} = \beta\alpha^{-1} Z\, W$ for some subword W of $\tilde{S}$. In particular, $W \neq \underline{x} \cdots$ and $W \neq \underline{y} \cdots$. Because $M(T)$ is assumed to have minimal length in its stable Auslander-Reiten component, the same is true for $M(\tilde{S})$ in its stable Auslander-Reiten component. Thus $W \neq 1_0$. Since we assume that $\tilde{S}$ is not the standard representative of a string of type $\beta\alpha^{-1}\lambda$, $W \neq \lambda$. Hence either (a) $W = \lambda\delta\, W'$, or (b) $W = \lambda\xi^{-1}\, W'$ for some subword W' of W. Analyzing these cases further, it follows that the stable endomorphism ring of every module in the stable Auslander-Reiten component containing $M(\tilde{S})$ has k-dimension at least 2. Hence it follows by Lemma 3.3 that this case cannot occur.

Therefore, $\tilde{S}$ in (3.2) must be the standard representative of a string of type $\beta\alpha^{-1}\lambda$. If $j = 0$, then $\theta^{-i_T}(\tilde{S}) = T$ and we define $S = \tilde{S}$. If $j = 1$, then by Remark 3.5,

$$\theta^{-i_T-1}(\Xi(\tilde{S^-})) = \theta^{-i_T-1}(\theta^2(\nu_2(\tilde{S}^{-1}))) = \theta^{-i_T+1}\left((\theta^2(\nu_1(\tilde{S})))^{-1}\right) = \left(\theta^{-i_T}(\nu_1(\tilde{S})\right)^{-1} = T^{-1}$$

and we define $S = \Xi(\tilde{S^-})$. By Definition 3.4, $\Xi(\tilde{S^-})$ is also the standard representative of a string of type $\beta\alpha^{-1}\lambda$. Hence in both cases $j = 0$ and $j = 1$, there exists $i \in \{0, 1, 2\}$ such that $\mathfrak{C}$ contains $M(\theta^i(S))$, where S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$. Since we have assumed that $\mathfrak{C}$ contains a module whose stable endomorphism ring is isomorphic to k, it follows by Lemma 3.6 that S satisfies condition (+). This completes the proof of Theorem 3.8. $\square$

Because of the prominent role played by standard representatives of strings S of type $\beta\alpha^{-1}\lambda$ in Theorem 3.8, the question arises which such S satisfy condition (+) from Lemma 3.6. We will give a precise answer to this question in Proposition 4.3.

We now look at the components of the stable Auslander-Reiten quiver of Λ_0 singled out by Theorem 3.8.

Proposition 3.9. *For $i \in \{0, 1, 2\}$, let $\mathfrak{C}_i$ be the component of the stable Auslander-Reiten quiver of Λ_0 containing the simple Λ_0-module $S_i = M(1_i)$. Define $(1_i)_h = \theta^i(\lambda\xi^{-1})$ and $(1_i)_{hh} = \theta^i(\lambda\xi^{-1}\delta\rho^{-1})$. The component $\mathfrak{C}_i$ is stable under Ω, and the modules in $\mathfrak{C}_i$ whose stable endomorphism rings are isomorphic to k are precisely the modules in the Ω-orbits of the modules $M(1_i)$, $M((1_i)_h)$ and $M((1_i)_{hh})$. Their universal deformation rings are*

$$R(\Lambda_0, M(1_i)) \cong k[[t]]/(t^2), \qquad R(\Lambda_0, M((1_i)_h)) \cong k, \qquad R(\Lambda_0, M((1_i)_{hh})) \cong k[[t]].$$

Proof. It suffices to consider $i = 0$. Using hooks and cohooks (see Definition 5.3), we see that $\mathfrak{C}_0$ is stable under Ω and that all Λ_0-modules in $\mathfrak{C}_0$ lie in the Ω-orbit of either

$$\begin{aligned} A_{n,0} &= M\left((\lambda\xi^{-1}\delta\rho^{-1}\beta\alpha^{-1})^n\right), \text{ or} \\ A_{n,1} &= M\left((\lambda\xi^{-1}\delta\rho^{-1}\beta\alpha^{-1})^n\lambda\xi^{-1}\right), \text{ or} \\ A_{n,2} &= M\left((\lambda\xi^{-1}\delta\rho^{-1}\beta\alpha^{-1})^n\lambda\xi^{-1}\delta\rho^{-1}\right) \end{aligned}$$

for some $n \geq 0$. Note that $M(1_0) = A_{0,0}$, $M((1_0)_h) = A_{0,1}$ and $M((1_0)_{hh}) = A_{0,2}$. Using Remark 5.4 and the description of the projective indecomposable Λ_0-modules in (3.1), it is straightforward to show that the stable endomorphism ring of $A_{0,j}$ is isomorphic to k for $j \in \{0, 1, 2\}$ and that $\mathrm{Ext}^1_{\Lambda_0}(A_{0,j}, A_{0,j})$ is isomorphic to k for $j \in \{0, 2\}$ and zero for $j = 1$. On the other hand, for $n \geq 1$, $A_{n,j}$ has a non-zero endomorphism which factors through $M(1_0)$ and which does not factor through a projective Λ_0-module.

Since $\mathrm{Ext}^1_{\Lambda_0}(A_{0,1}, A_{0,1}) = 0$, it follows that $R(\Lambda_0, A_{0,1}) \cong k$. Since $\mathrm{Ext}^1_{\Lambda_0}(A_{0,j}, A_{0,j})$ is isomorphic to k for $j \in \{0, 2\}$, it follows that $R(\Lambda_0, A_{0,j})$ is a quotient of $k[[t]]$ for $j \in \{0, 2\}$.

The module $A_{0,0}$ lies in a non-split short exact sequence of Λ_0-modules

$$0 \to A_{0,0} \to M(\alpha) \to A_{0,0} \to 0. \tag{3.3}$$

Therefore, $M(\alpha)$ defines a non-trivial lift of $A_{0,0}$ over $k[[t]]/(t^2)$. This implies that there is a unique surjective k-algebra homomorphism $\psi : R(\Lambda_0, A_{0,0}) \to k[[t]]/(t^2)$ in $\hat{\mathcal{C}}$ corresponding to the deformation defined by $M(\alpha)$. We need to show that ψ is an isomorphism. Suppose this is false.

Then there exists a surjective k-algebra homomorphism $\psi_0 : R(\Lambda_0, A_{0,0}) \to k[[t]]/(t^3)$ in $\hat{\mathcal{C}}$ such that $\pi \circ \psi_0 = \psi$ where $\pi : k[[t]]/(t^3) \to k[[t]]/(t^2)$ is the natural projection. Let M_0 be a $k[[t]]/(t^3) \otimes_k \Lambda_0$-module which defines a lift of $A_{0,0}$ over $k[[t]]/(t^3)$ corresponding to ψ_0. Because $M_0/t^2M_0 \cong M(\alpha)$ and $t^2M_0 \cong A_{0,0}$, we obtain a non-split short exact sequence of $k[[t]]/(t^3) \otimes_k \Lambda_0$-modules

$$0 \to A_{0,0} \to M_0 \to M(\alpha) \to 0. \tag{3.4}$$

Since $\mathrm{Ext}^1_{\Lambda_0}(M(\alpha), A_{0,0}) = 0$, this sequence splits as a sequence of Λ_0-modules. Hence $M_0 = A_{0,0} \oplus M(\alpha)$ as Λ_0-modules. Writing elements of M_0 as (a, m) where $a \in A_{0,0}$ and $m \in M(\alpha)$, the t-action on M_0 is given as $t\,(a, m) = (\sigma(m), t\, m)$, where $\sigma : M(\alpha) \to A_{0,0}$ is a surjective Λ_0-module homomorphism. Since for all such σ we have $\sigma(t\, m) = 0 = t^2\, m$, it follows that $t^2\,(a, m) = (\sigma(t\, m), t^2\, m) = (0, 0)$ for all $a \in A_{0,0}$ and $m \in M(\alpha)$. But this is a contradiction to $t^2M_0 \cong A_{0,0}$. Thus ψ is a k-algebra isomorphism and $R(\Lambda_0, A_{0,0}) \cong k[[t]]/(t^2)$.

Let $\underline{a} = \lambda\xi^{-1}\delta\rho^{-1}$ and consider $A_{0,2} = M(\underline{a})$. Then $A_{0,2}$ lies in a non-split short exact sequence of Λ_0-modules

$$0 \to A_{0,2} \to M(\underline{a}\beta\underline{a}) \to A_{0,2} \to 0. \tag{3.5}$$

Therefore, $M(\underline{a}\beta\underline{a})$ defines a non-trivial lift of $A_{0,2}$ over $k[[t]]/(t^2)$. Let $\{b_0, b_1, \ldots, b_4\}$ be a canonical k-basis of $A_{0,2}$ relative to the representative $\underline{a}$, and let for each arrow $\zeta \in \{\alpha, \rho, \xi, \beta, \delta, \lambda\}$, X_ζ be the 5×5 matrix with entries in k describing the action of ζ on $A_{0,2}$ with respect to $\{b_0, b_1, \ldots, b_4\}$ (see Definition 5.1). Let $L_{0,2}$ be a free $k[[t]]$-module of rank 5, and let $\{B_0, B_1, \ldots, B_4\}$ be a basis of $L_{0,2}$ over $k[[t]]$. Viewing k as a subalgebra of $k[[t]]$, define a Λ_0-module structure on $L_{0,2}$ by letting $\zeta \in \{\alpha, \rho, \xi, \delta, \lambda\}$ act as the matrix X_ζ and β as the matrix $X_\beta + tE_{4,0}$ with respect to the basis $\{B_0, B_1, \ldots, B_4\}$. Here $E_{4,0}$ is the elementary 5×5 matrix which sends B_0 to B_4 and all other basis elements to 0. Then $L_{0,2}$ is a $k[[t]] \otimes_k \Lambda_0$-module which is free over $k[[t]]$ and $L_{0,2}/tL_{0,2} \cong A_{0,2}$. Hence $L_{0,2}$ defines a lift of $A_{0,2}$ over $k[[t]]$, and there exists a unique k-algebra homomorphism $\varphi : R(\Lambda_0, A_{0,2}) \to k[[t]]$ in $\hat{\mathcal{C}}$ corresponding to the deformation defined by $L_{0,2}$. Because $L_{0,2}/t^2L_{0,2} \cong M(\underline{a}\beta\underline{a})$ as Λ_0-modules, $L_{0,2}/t^2L_{0,2}$ defines a non-trivial lift of $A_{0,2}$ over $k[[t]]/(t^2)$. Thus φ is surjective. Since $R(\Lambda_0, A_{0,2})$ is a quotient of $k[[t]]$, this implies that φ is an isomorphism, and hence $R(\Lambda_0, A_{0,2}) \cong k[[t]]$. This proves Proposition 3.9. □

Proposition 3.10. *Let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$ satisfying condition (+) from Lemma 3.6. Then ${}_hS = \lambda\delta\rho^{-1}S$ and ${}_{hh}S = \delta\beta\alpha^{-1}({}_hS)$. For $i \in \{0, 1, 2\}$, let $\mathfrak{C}_{S,i}$ be the component of the stable Auslander-Reiten quiver of Λ_0 containing $M(\theta^i(S))$. The component $\mathfrak{C}_{S,i}$ is stable under Ω, and the modules in $\mathfrak{C}_{S,i}$ whose stable endomorphism rings are isomorphic to k are precisely the modules in the Ω-orbits of the modules $M(\theta^i(S))$, $M(\theta^i({}_hS))$ and $M(\theta^i({}_{hh}S))$. Their universal deformation rings are*

$$R(\Lambda_0, M(\theta^i(S))) \cong k[[t]]/(t^2), \qquad R(\Lambda_0, M(\theta^i({}_hS))) \cong k, \qquad R(\Lambda_0, M(\theta^i({}_{hh}S))) \cong k[[t]].$$

Proof. It suffices to consider the case when $i = 0$. Using hooks and cohooks (see Definition 5.3), we see that $\mathfrak{C}_{S,0}$ is stable under Ω and that all Λ_0-modules in $\mathfrak{C}_{S,0}$ lie in the Ω-orbit of either

$$\begin{aligned} V_{n,0} &= M\left(S(\delta\rho^{-1}\beta\alpha^{-1}\lambda\xi^{-1})^n\delta\rho^{-1}\right), \text{ or} \\ V_{n,1} &= M\left(S(\delta\rho^{-1}\beta\alpha^{-1}\lambda\xi^{-1})^n\delta\rho^{-1}\beta\alpha^{-1}\right), \text{ or} \\ V_{n,2} &= M\left(S(\delta\rho^{-1}\beta\alpha^{-1}\lambda\xi^{-1})^{n+1}\right) \end{aligned}$$

for some $n \geq 0$. Since $S = S^-$ when S satisfies condition (+), we have $M(S) \cong \Omega(V_{0,0})$, $M({}_hS) \cong \Omega(V_{0,1})$ and $M({}_{hh}S) \cong \Omega(V_{0,2})$. By Lemma 3.6, the stable endomorphism ring of $M(S)$ is isomorphic to k. Using Remark 5.4 together with similar arguments as in the proof of Lemma 3.6, it follows that the stable endomorphism ring of $V_{0,j}$ is isomorphic to k for $j \in \{0, 1, 2\}$. Similarly, we analyze $\mathrm{Ext}^1_{\Lambda_0}(V_{0,j}, V_{0,j})$ to see that it is isomorphic to k for $j \in \{0, 2\}$ and zero for $j = 1$. On the other hand, for $n \geq 1$, $V_{n,j}$ has a non-zero endomorphism which factors through $M(1_1)$ and which does not factor through a projective Λ_0-module.

The universal deformation rings of $V_{0,0}$, $V_{0,1}$ and $V_{0,2}$ are determined in a similar way as in the proof of Proposition 3.9. It follows that $R(\Lambda_0, V_{0,0}) \cong k[[t]]/(t^2)$, $R(\Lambda_0, V_{0,1}) \cong k$ and $R(\Lambda_0, V_{0,2}) \cong k[[t]]$. Using Theorem 2.6(iv), this proves Proposition 3.10. $\square$

Proposition 3.11. *Let $\mathfrak{T}_1$ and $\mathfrak{T}_2$ be the two 3-tubes of the stable Auslander-Reiten quiver of Λ_0. Then $\Omega(\mathfrak{T}_1) = \mathfrak{T}_2$. Define $T = \alpha^{-1}$. Then $T_h = \alpha^{-1}\lambda\xi^{-1}$ and $T_{hh} = \alpha^{-1}\lambda\xi^{-1}\delta\rho^{-1}$. The modules in $\mathfrak{T}_1 \cup \mathfrak{T}_2$ whose stable endomorphism rings are isomorphic to k are precisely the modules in the Ω-orbits of the modules $M(T)$, $M(T_h)$ and $M(T_{hh})$. Their universal deformation rings are*

$$R(\Lambda_0, M(T)) \cong k \cong R(\Lambda_0, M(T_h)), \qquad R(\Lambda_0, M(T_{hh})) \cong k[[t]].$$

Proof. Using the description of the projective indecomposable Λ_0-modules in (3.1), we see that $\Omega(\mathfrak{T}_1) = \mathfrak{T}_2$. Using hooks and cohooks (see Definition 5.3) and Remark 5.4, it is straightforward to show that the only Λ_0-modules in $\mathfrak{T}_1 \cup \mathfrak{T}_2$ whose stable endomorphism rings are isomorphic to k lie in the Ω-orbit of either $C_0 = M(T)$, $C_1 = M(T_h)$ or $C_2 = M(T_{hh})$. Since $\mathrm{Ext}^1_{\Lambda_0}(C_j, C_j) = 0$ for $j \in \{0,1\}$, we have $R(\Lambda_0, C_j) \cong k$ for $j \in \{0,1\}$. Since $\mathrm{Ext}^1_{\Lambda_0}(C_2, C_2) \cong k$, it follows that $R(\Lambda_0, C_2)$ is a quotient of $k[[t]]$. Using similar arguments as in the proof of Proposition 3.9, we obtain that $R(\Lambda_0, C_2) \cong k[[t]]$, which proves Proposition 3.11. $\square$

3.2. Stable endomorphism rings and universal deformation rings of band modules for Λ_0.

In this section, we describe all band modules for Λ_0 whose stable endomorphism rings are isomorphic to k and determine their universal deformation rings.

It follows from [15] that if B is a band, $\mu \in k^*$ and $n \geq 2$ is an integer, then $\underline{\mathrm{End}}_{\Lambda_0}(M(B, \mu, n))$ has k-dimension at least 2. Hence we can concentrate on the band modules $M_{B,\mu} = M(B, \mu, 1)$.

For the description of all the band modules whose stable endomorphism rings are isomorphic to k, we need to look at special bands, which we call bands of type 0 and which are defined using the words of type 0 and their rotations of type 0 introduced in Definition 3.2.

Definition 3.12. (i) We say a word B representing a band is a *top-socle representative* if

$$B = C_1 C_2^{-1} C_3 C_4^{-1} \cdots C_{2l-1} C_{2l}^{-1}$$

where $l \geq 1$, $C_{2i-1} \in \{\beta, \beta\lambda, \delta, \delta\beta, \lambda, \lambda\delta\}$ and $C_{2i} \in \{\alpha, \rho, \xi\}$ for $1 \leq i \leq l$. The words $C_1, C_2, \ldots, C_{2l}$ are called the *top-socle pieces* of B. The positive integer l is called the *top-socle length* of B. For $1 \leq j \leq 2l$, $s(C_j)$ is called the *top* of C_j and $e(C_j)$ is called the *socle* of C_j. The collection of all the tops (resp. of all the socles) of the C_{2i-1}, $1 \leq i \leq l$, where we count multiplicities, is called the *top* (resp. the *socle*) of B.

(ii) We say a band is a *band of type* 0 if it has a top-socle representative B which is a word of type 0, as defined in Definition 3.2(ii). In other words,

$$B = \underline{z}_0\, \underline{z}_1 \, \cdots \, \underline{z}_{n-1}$$

where $n \geq 1$ and $\underline{z}_0, \ldots, \underline{z}_{n-1} \in \{\underline{x}, \underline{y}\}$. A *standard representative of a band of type* 0 is a top-socle representative which is a word of type 0. This is unique up to rotations of type 0, as defined in Definition 3.2(iv).

(iii) Let $B = \underline{z}_0\, \underline{z}_1 \cdots \underline{z}_{n-1}$ be a standard representative of a band of type 0. A *rotated subpattern* U of B is a subpattern of $\rho_i^{(0)}(B)$ for some $0 \leq i \leq n-1$, where $\rho_i^{(0)}$ is as in Definition 3.2(iv). Thus either $U = 1_0$ or $U = \underline{z}_i\, \underline{z}_{i+1} \cdots \underline{z}_j$ for some $0 \leq i \leq j \leq n-1$ or $U = \underline{z}_i\, \underline{z}_{i+1} \cdots \underline{z}_{n-1} \underline{z}_0 \cdots \underline{z}_j$ for some $0 \leq j < i \leq n-1$. The *inverse pattern* U^- of U is given as in Definition 3.2(iii).

(iv) Let $B = \underline{z}_0\, \underline{z}_1 \cdots \underline{z}_{n-1}$ be a standard representative of a band of type 0, and let $\omega \in \{0, 1, \ldots, n-1\}$. We say B *allows a wrap-around at* ω if $\underline{z}_i = \underline{z}_{\omega - i}$ for all i, where the indices are taken modulo n.

Remark 3.13. Suppose $B = \underline{z}_0\, \underline{z}_1 \cdots \underline{z}_{n-1}$ is a standard representative of a band of type 0.

(i) Since B is not a power of a smaller word, B allows at most one wrap-around.

(ii) Suppose B allows a wrap-around at ω. Let $\mu,\tilde{\mu}\in k^*$, and let $M_{B,\mu}=M(B,\mu,1)$ and $M_{B,\tilde{\mu}}=M(B,\tilde{\mu},1)$. We obtain the following sequence of canonical homomorphisms from $M_{B,\mu}$ to $M_{B,\tilde{\mu}}$, as defined in Remark 5.5, associated to the wrap-around at ω.

Let $\{b_0,b_1,\ldots,b_{5n-1}\}$ (resp. $\{\tilde{b}_0,\tilde{b}_1,\ldots,\tilde{b}_{5n-1}\}$) be a canonical k-basis for $M_{B,\mu}$ (resp. $M_{B,\tilde{\mu}}$) relative to the representative B (see Definition 5.2). Let $0\le j\le n-1$. Define $\nu_j=1$ and $\epsilon_j=2$ if $\underline{z}_j=\underline{x}$ (resp. $\nu_j=2$ and $\epsilon_j=1$ if $\underline{z}_j=\underline{y}$). Then $\xi_{j,1}$ (resp. $\xi_{j,2}$) with $\xi_{j,1}(b_{5j+\nu_j})=\tilde{b}_{5(\omega-j)+\nu_j+1}$ and $\xi_{j,1}(b_i)=0$ for all other i (resp. $\xi_{j,2}(b_{5j+4})=\tilde{b}_{5(\omega-j)}$ and $\xi_{j,2}(b_i)=0$ for all other i) defines a canonical homomorphism from $M_{B,\mu}$ to $M_{B,\tilde{\mu}}$ of string type 1_{ϵ_j} (resp. 1_0).

Considering the projective indecomposable Λ_0-modules, it follows that $\xi_{j,1}+\xi_{j,2}$ for $j\in\{0,1,\ldots,n-1\}-\{\omega\}$ and $\xi_{\omega,1}+\tilde{\mu}\xi_{\omega,2}$ all factor through projective Λ_0-modules. Also, $\xi_{j,2}+\xi_{j+1,1}$ for $j\in\{0,1,\ldots,n-2\}$ and $\xi_{n-1,2}+\mu\xi_{0,1}$ all factor through projective Λ_0-modules. We conclude that every homomorphism in $\{\xi_{j,1},\xi_{j,2}\mid 0\le j\le n-1\}$ factors through a projective Λ_0-module if and only if $\mu\tilde{\mu}\neq 1$.

Lemma 3.14. *Let $B=\underline{z}_0\underline{z}_1\cdots\underline{z}_{n-1}$ be a standard representative of a band B of type 0 as in Definition 3.12, let $\mu\in k^*$ and let $M_{B,\mu}=M(B,\mu,1)$. The stable endomorphism ring of $M_{B,\mu}$, $\underline{\mathrm{End}}_{\Lambda_0}(M_{B,\mu})$, is isomorphic to k if and only if $\mu\neq\pm1$ and B has the following property:*

(++) *If there are rotated subpatterns U,V of B such that $B\sim_{r,(0)}\underline{x}\,U\,\underline{x}\,V$, then for all rotated subpatterns W of B, we have $B\not\sim_{r,(0)}\underline{y}\,U\,\underline{y}\,W$.*

Here $\sim_{r,(0)}$ is the equivalence relation introduced in Definition 3.2(iv).

Proof. Since for $B\in\{\underline{x},\underline{y}\}$, $\underline{\mathrm{End}}_{\Lambda_0}(M_{B,\mu})\cong k$ if and only if $\mu\neq\pm1$ and since $\underline{x}$ and $\underline{y}$ have property (++), Lemma 3.14 follows for $n=1$. For the remainder of the proof, let $B=\underline{z}_0\underline{z}_1\cdots\underline{z}_{n-1}$ be a standard representative of a band of type 0 with $n\ge2$, and let $\mu\in k^*$. We analyze the different possibilities for $M_{B,\mu}$ such that $\underline{\mathrm{End}}_{\Lambda_0}(M_{B,\mu})$ has k-dimension at least 2.

Suppose first that there exists a string T of length at least 2 such that $M_{B,\mu}$ has a canonical endomorphism α_T of string type T, as described in Remark 5.5. In particular, $\underline{\mathrm{End}}_{\Lambda_0}(M_{B,\mu})$ has k-dimension at least 2. It follows from the definition of bands of type 0 that

$$T\sim_s\beta\alpha^{-1}U\lambda$$

for some rotated subpattern U of B and that there exist rotated subpatterns V,W of B such that

(‡) $B\sim_{r,(0)}\underline{x}\,U\,\underline{x}\,V$ and $B\sim_{r,(0)}\underline{y}\,U\,\underline{y}\,W$.

On the other hand, if B satisfies (‡), then $M_{B,\mu}$ has a canonical endomorphism factoring through a string module with at least three composition factors.

Next let T be a string of length 0 or 1 such that $M_{B,\mu}$ has a canonical endomorphism α_T of string type T. Then either $T=\delta$ or $T=1_u$ for some $u\in\{0,1,2\}$.

Analyzing these cases further, it follows that if $T=\delta$ then α_T does not factor through a projective Λ_0-module only if B satisfies (‡) for certain rotated subpatterns U,V,W of B.

Next let $T=1_u$ for some $u\in\{0,1,2\}$. Then α_T does not factor through a projective Λ_0-module only if either (a) B satisfies (‡) for certain U,V,W, or (b) B has property (++) and $\mu=\pm1$. To explain (b), we first note that if B has property (++), then B allows a wrap-around at ω for a unique $\omega\in\{0,1,\ldots,n-1\}$ (see Proposition 4.4). Using Remark 3.13(ii), it follows that if B satisfies (++) then α_T does not factor through a projective Λ_0-module if and only if there exist $0\le j\le n-1$ and $s\in\{1,2\}$ such that $\alpha_T=\xi_{j,s}$ and $\mu=\pm1$. This completes the proof of Lemma 3.14. □

Definition 3.15. Define

$$\underline{p}=\lambda\xi^{-1}\delta\rho^{-1}\beta\alpha^{-1}\quad\text{and}\quad\underline{q}=\lambda\delta\rho^{-1}\beta\lambda\xi^{-1}\delta\beta\alpha^{-1}.\tag{3.6}$$

Theorem 3.16. *Let M be a band module for Λ_0. Then the stable endomorphism ring of M is isomorphic to k if and only if either*

(i) *there exist $\mu\in k^*$ and $B\in\{\underline{p},\underline{q}\}$ such that $M\cong M_{B,\mu}$, or*

(ii) *there exist* $i \in \{0,1,2\}$, $\mu \in k^*$ *with* $\mu \neq \pm 1$, *and a standard representative* B *of a band of type* 0 *satisfying condition* $(++)$ *from Lemma 3.14 such that* $M \cong M_{\theta^i(B),\mu}$.

Proof. Let C be a top-socle representative of a band whose top-socle length is $l \geq 1$, and let $\mu \in k^*$ be such that $M \cong M_{C,\mu}$. As in Definition 3.12(i), C has the form

$$C = D_1 D_2^{-1} D_3 D_4^{-1} \cdots D_{2l-1} D_{2l}^{-1}$$

where $D_{2j-1} \in \{\beta, \beta\lambda, \delta, \delta\beta, \lambda, \lambda\delta\}$ and $D_{2j} \in \{\alpha, \rho, \xi\}$. In particular, $l \geq 2$.

If all D_{2j-1} have the same length, then either $C \sim_r \underline{p}$ or $C \sim_r \underline{q}$. In this case the stable endomorphism ring of $M_{C,\mu}$ is isomorphic to k for all μ.

For the remainder of the proof suppose that not all D_{2j-1} have the same length. Then there exists $i_0 \in \{0,1,2\}$ such that C has a subword of the form $\theta^{i_0}(\underline{x})$ or $\theta^{i_0}(\underline{y})$. This means that we can write

$$\tilde{C} \overset{\text{def}}{=} \theta^{-i_0}(C) = Z_0 Z_1 \cdots Z_{m-1}$$

where for all $0 \leq j \leq m-1$, Z_j is a sequence of top-socle pieces of $\tilde{C}$ such that $Z_j = \lambda \cdots \alpha^{-1}$ and 1_0 occurs exactly once in the top of Z_j. Hence $Z_j \in \{\underline{p}_r, \underline{q}_r, \underline{x}_r, \underline{y}_r \mid r \geq 0\}$, where

$$\underline{p}_r = \lambda\xi^{-1}\delta\rho^{-1}\left(\beta\lambda\xi^{-1}\delta\rho^{-1}\right)^r \beta\alpha^{-1}, \quad \underline{q}_r = \lambda\left(\delta\rho^{-1}\beta\lambda\xi^{-1}\right)^{r+1}\delta\beta\alpha^{-1},$$
$$\underline{x}_r = \lambda\xi^{-1}\left(\delta\rho^{-1}\beta\lambda\xi^{-1}\right)^r \delta\beta\alpha^{-1}, \quad \underline{y}_r = \lambda\delta\rho^{-1}\left(\beta\lambda\xi^{-1}\delta\rho^{-1}\right)^r \beta\alpha^{-1}.$$

In particular, $\underline{p}_0 = \underline{p}$, $\underline{q}_0 = \underline{q}$, $\underline{x}_0 = \underline{x}$, $\underline{y}_0 = \underline{y}$. Using that at least one of the Z_j is $\underline{x}_0$ or $\underline{y}_0$, we obtain that the stable endomorphism ring of $M_{\tilde{C},\mu}$, and hence of $M_{C,\mu}$, has k-dimension at least 2 for all $\mu \in k^*$ except in the following cases:

(a) $\tilde{C}$ only contains $\underline{x}_0$ and $\underline{y}_0$; or
(b) $\tilde{C}$ only contains $\underline{x}_0$ and $\underline{x}_1$; or
(c) $\tilde{C}$ only contains $\underline{y}_0$ and $\underline{y}_1$.

Suppose that $\tilde{C}$ is as in one of the cases (a) – (c). Then there exists $i_1 \in \{0,1,2\}$ such that a rotation of $\theta^{i_1}(\tilde{C})$ is a standard representative of a band of type 0. Thus there exist $i \in \{0,1,2\}$ and a standard representative B of a band of type 0 such that $M_{C,\mu} \cong M_{\theta^i(B),\mu}$. By Lemma 3.14, it follows that the stable endomorphism ring of $M_{C,\mu}$ is isomorphic to k if and only if $\mu \neq \pm 1$ and B satisfies condition $(++)$. This completes the proof of Theorem 3.16. □

Because of the prominent role played by standard representatives of bands B of type 0 in Theorem 3.16, the question arises which such B satisfy condition $(++)$ from Lemma 3.14. We will give a precise answer to this question in Proposition 4.4.

We now look at the band modules for Λ_0 singled out by Theorem 3.16.

Proposition 3.17. *Let* B *be a top-socle representative of a band and let* $\mu \in k^*$.

(i) *If* $\mu \in k^*$ *and* $\{B, B'\} = \{\underline{p}, \underline{q}\}$, *then* $\Omega(M_{B,\mu}) \cong M_{B',-\mu^{-1}}$.
(ii) *If* $\mu \in k^* - \{\pm 1\}$ *and there exists* $i \in \{0,1,2\}$ *such that* $\theta^{-i}(B)$ *is a standard representative of a band of type* 0 *satisfying condition* $(++)$ *from Lemma 3.14, then* $\Omega(M_{B,\mu}) \cong M_{B,\mu^{-1}}$.

In each of the above cases, the universal deformation ring is $R(\Lambda_0, M_{B,\mu}) \cong k[[t]]$.

Proof. Using that $\Lambda_0 = P_2 \oplus P_1 \oplus P_0$ is a projective Λ_0-module cover of $M_{\underline{p},\mu}$ and analyzing the kernel $\Omega(M_{\underline{p},\mu})$ of the natural surjection $\Lambda_0 \to M_{\underline{p},\mu}$, it follows that $\Omega(M_{\underline{p},\mu}) \cong M_{\underline{q},-\mu^{-1}}$. Since $\Omega^2(M_{\underline{p},\mu}) \cong M_{\underline{p},\mu}$, we also have $\Omega(M_{\underline{q},\mu}) \cong M_{\underline{p},-\mu^{-1}}$. Using Remark 5.5, we see that the vector space $\mathrm{Hom}_{\Lambda_0}(M_{\underline{q},-\mu^{-1}}, M_{\underline{p},\mu})$ is 3-dimensional over k, but the subspace of homomorphisms factoring through projective Λ_0-modules has k-dimension 2. Hence

$$\mathrm{Ext}^1_{\Lambda_0}(M_{\underline{p},\mu}, M_{\underline{p},\mu}) \cong \underline{\mathrm{Hom}}_{\Lambda_0}(\Omega(M_{\underline{p},\mu}), M_{\underline{p},\mu}) \cong \underline{\mathrm{Hom}}_{\Lambda_0}(M_{\underline{q},-\mu^{-1}}, M_{\underline{p},\mu}) \cong k.$$

This implies that $R(\Lambda_0, M_{\underline{p},\mu})$ is a quotient of $k[[t]]$. Using similar arguments as in the proof of Proposition 3.9, we obtain that $R(\Lambda_0, M_{\underline{p},\mu}) \cong k[[t]]$. By Theorem 2.6(iv), we also get $R(\Lambda_0, M_{\underline{q},\mu}) \cong k[[t]]$, which proves part (i).

Suppose now that μ, i and B are as in part (ii). Without loss of generality, we may assume that $i = 0$, which means that B is a standard representative of a band of type 0 satisfying condition (++). Then B allows a unique wrap-around (see Proposition 4.4), which implies $B \sim_{r,(0)} B^-$. Since the top-socle length of B is even, it follows that $\Omega(M_{B,\mu}) \cong M_{B,\mu^{-1}}$. Using Remark 5.5, we see that the k-dimension of the vector space $\mathrm{Hom}_{\Lambda_0}(M_{B,\mu^{-1}}, M_{B,\mu})$ is one less than the k-dimension of $\mathrm{End}_{\Lambda_0}(M_{B,\mu})$, since there is no identity homomorphism in the former space. Arguing in a similar way as in the proof that the stable endomorphism ring of $M_{B,\mu}$ is isomorphic to k, we see that all the canonical homomorphisms $M_{B,\mu^{-1}} \to M_{B,\mu}$ factor through projective Λ_0-modules except possibly the homomorphisms $\xi_{j,1}, \xi_{j,2}$, $0 \le j \le n-1$, from Remark 3.13(ii). Since $\mu^{-1}\mu = 1$, it follows that

$$\mathrm{Ext}^1_{\Lambda_0}(M_{B,\mu}, M_{B,\mu}) \cong \underline{\mathrm{Hom}}_{\Lambda_0}(\Omega(M_{B,\mu}), M_{B,\mu}) \cong \underline{\mathrm{Hom}}_{\Lambda_0}(M_{B,\mu^{-1}}, M_{B,\mu}) \cong k.$$

This implies that $R(\Lambda_0, M_{B,\mu})$ is a quotient of $k[[t]]$. Using similar arguments as in the proof of Proposition 3.9, we obtain that $R(\Lambda_0, M_{B,\mu}) \cong k[[t]]$. This completes the proof of Proposition 3.17 □

4. Appendix: Strings and bands for Λ_0 satisfying (+) and (++) from Section 3

We assume the notation from §3. In particular, k is an algebraically closed field of arbitrary characteristic and Λ_0 is the basic k-algebra $\Lambda_0 = kQ/I$ where Q and I are as in Figure 1. Because of the prominent role played by standard representatives of strings S of type $\beta\alpha^{-1}\lambda$ satisfying condition (+) in Theorem 3.8, respectively by standard representatives of bands B of type 0 satisfying condition (++) in Theorem 3.16, the question arises if one can give an explicit description of such S and B. In this appendix, we give a precise answer to this question in Propositions 4.3 and 4.4.

Considering words of type 0 of small type-0-length suggests that to answer this question one needs to use an inductive process. The key ingredients in this inductive process are described in the following definition. We assume Definition 3.2.

Definition 4.1. (i) Let ℓ be a positive integer, let $(i_1, \ldots, i_{\ell-1})$ be an $(\ell-1)$-tuple with $i_j \in \{0,1\} \mod 2$ for $1 \le j \le \ell-1$, and let $(a_1, \ldots, a_{\ell-1})$ be an $(\ell-1)$-tuple of positive integers. We define

$$N_0 = \underline{x}, \qquad N_1 = \underline{y};$$

and for $1 \le j \le \ell - 1$,

$$\begin{aligned} N_{i_1,\ldots,i_j,0}(a_1,\ldots,a_j) &= N_{i_1,\ldots,i_j}(a_1,\ldots,a_{j-1})^{a_j}\, N_{i_1,\ldots,i_{j-1},(i_j)+1}(a_1,\ldots,a_{j-1}), \\ N_{i_1,\ldots,i_j,1}(a_1,\ldots,a_j) &= N_{i_1,\ldots,i_j}(a_1,\ldots,a_{j-1})^{a_j+1}\, N_{i_1,\ldots,i_{j-1},(i_j)+1}(a_1,\ldots,a_{j-1}). \end{aligned}$$

(ii) Let ℓ be a positive integer, let $(i_1, \ldots, i_{\ell-1})$ be an $(\ell-1)$-tuple with $i_j \in \{0,1\} \mod 2$ for $1 \le j \le \ell-1$, and let $(a_1, \ldots, a_{\ell-1})$ be an $(\ell-1)$-tuple of positive integers. Define

$$\begin{aligned} \underline{x}^{(\ell)} &= N_{i_1,\ldots,i_{\ell-1},0}(a_1,\ldots,a_{\ell-1}), \\ \underline{y}^{(\ell)} &= N_{i_1,\ldots,i_{\ell-1},1}(a_1,\ldots,a_{\ell-1}). \end{aligned}$$

We say a word Z is a *word of type* 0 *on level* ℓ if $Z = 1_0$ or

$$Z = \underline{z}_0^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$$

where $n \ge 1$ and $\underline{z}_0^{(\ell)}, \ldots, \underline{z}_{n-1}^{(\ell)} \in \{\underline{x}^{(\ell)}, \underline{y}^{(\ell)}\}$. We call n the *type*-0-*length on level* ℓ of Z, where $n = 0$ corresponds to $Z = 1_0$. Note that for $\ell = 1$, this is the original definition of a word of type 0 from Definition 3.2.

If we want to emphasize the tuples used to define $\underline{x}^{(\ell)}$ and $\underline{y}^{(\ell)}$, we say Z is a word of type 0 on level ℓ relative to the $(\ell-1)$-tuples $(i_1, \ldots, i_{\ell-1})$ and $(a_1, \ldots, a_{\ell-1})$.

(iii) Let $Z = \underline{z}_0^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$ be a word of type 0 on level ℓ. A *subpattern* U of Z *on level* ℓ is a subword of Z such that either $U = 1_0$ or $U = \underline{z}_i^{(\ell)} \underline{z}_{i+1}^{(\ell)} \cdots \underline{z}_j^{(\ell)}$ for some $0 \le i \le j \le n-1$.

(iv) Let $Z = \underline{z}_0^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$ be a word of type 0 on level ℓ with $n \geq 1$. For $0 \leq i \leq n-1$, we define the i^{th} *rotation of* Z *of type* 0 *on level* ℓ to be

$$\rho_i^{(0,\ell)}(Z) = \underline{z}_i^{(\ell)} \underline{z}_{i+1}^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)} \underline{z}_0^{(\ell)} \cdots \underline{z}_{i-1}^{(\ell)}.$$

Define $\sim_{r,(0,\ell)}$ to be the equivalence relation on all words of type 0 on level ℓ of type-0-length on level ℓ at least 1 such that $Z \sim_{r,(0,\ell)} Z'$ if and only if $Z = \rho_i^{(0,\ell)}(Z')$ for some i. Note that for $\ell = 1$, the equivalence relation $\sim_{r,(0,\ell)}$ coincides with the equivalence relation $\sim_{r,(0)}$ from Definition 3.2.

(v) Let ℓ be a positive integer, let $(i_1, \dots, i_{\ell-1})$ be an $(\ell-1)$-tuple with $i_j \in \{0,1\} \mod 2$ for $1 \leq j \leq \ell - 1$, and let $(a_1, \dots, a_{\ell-1})$ be an $(\ell-1)$-tuple of positive integers. Define $\underline{x}^{(\ell)}$ and $\underline{y}^{(\ell)}$ as in part (ii) and define

$$\begin{aligned} \underline{b}^{(\ell)} &= \beta\alpha^{-1} N_{i_1,0}(a_1)\, N_{i_1,i_2,0}(a_1,a_2) \cdots N_{i_1,\dots,i_{\ell-1},0}(a_1,\dots,a_{\ell-1}), \\ \underline{l}^{(\ell)} &= N_{i_1,\dots,i_{\ell-1}}(a_1,\dots,a_{\ell-2})^{a_{\ell-1}} \cdots N_{i_1,i_2}(a_1)^{a_2}\, (N_{i_1})^{a_1}\, \lambda. \end{aligned}$$

We say a word S is the *standard representative of a string of type* $\beta\alpha^{-1}\lambda$ *on level* ℓ if

$$S = \underline{b}^{(\ell)} Z\, \underline{l}^{(\ell)}$$

where Z is a word of type 0 on level ℓ. Note that for $\ell = 1$, this is the original definition of the standard representative of a string of type $\beta\alpha^{-1}\lambda$ from Definition 3.4.

If we want to emphasize the tuples used to define $\underline{x}^{(\ell)}$, $\underline{y}^{(\ell)}$, $\underline{b}^{(\ell)}$ and $\underline{l}^{(\ell)}$, we say S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ relative to the $(\ell-1)$-tuples $(i_1, \dots, i_{\ell-1})$ and $(a_1, \dots, a_{\ell-1})$.

A *subpattern* U of S *on level* ℓ is a subpattern of Z on level ℓ, as defined in part (iii).

(vi) Let ℓ be a positive integer, let $(i_1, \dots, i_{\ell-1})$ be an $(\ell-1)$-tuple with $i_j \in \{0,1\} \mod 2$ for $1 \leq j \leq \ell - 1$, and let $(a_1, \dots, a_{\ell-1})$ be an $(\ell-1)$-tuple of positive integers. Define $\underline{x}^{(\ell)}$ and $\underline{y}^{(\ell)}$ as in part (ii). We say a band is a *band of type* 0 *on level* ℓ if it has a top-socle representative B which is a word of type 0 on level ℓ. In other words,

$$B = \underline{z}_0^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$$

where $n \geq 1$ and $\underline{z}_0^{(\ell)}, \dots, \underline{z}_{n-1}^{(\ell)} \in \{\underline{x}^{(\ell)}, \underline{y}^{(\ell)}\}$. A *standard representative of a band of type* 0 *on level* ℓ is a top-socle representative which is a word of type 0 on level ℓ. This is unique up to rotations of type 0 on level ℓ, as defined in part (iv). Note that for $\ell = 1$, this is the original definition of a band of type 0 and a standard representative of a band of type 0 from Definition 3.12.

If we want to emphasize the tuples used to define $\underline{x}^{(\ell)}$ and $\underline{y}^{(\ell)}$, we say B is a standard representative of a band of type 0 on level ℓ relative to the $(\ell-1)$-tuples $(i_1, \dots, i_{\ell-1})$ and $(a_1, \dots, a_{\ell-1})$.

(vii) Let $B = \underline{z}_0^{(\ell)}\, \underline{z}_1^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$ be a standard representative of a band of type 0 on level ℓ. A *rotated subpattern* U of B *on level* ℓ is a subpattern of $\rho_i^{(0,\ell)}(B)$ for some $0 \leq i \leq n-1$, where $\rho_i^{(0,\ell)}$ is as in part (iv). Thus either $U = 1_0$ or $U = \underline{z}_i^{(\ell)}\, \underline{z}_{i+1}^{(\ell)} \cdots \underline{z}_j^{(\ell)}$ for some $0 \leq i \leq j \leq n$ or $U = \underline{z}_i^{(\ell)}\, \underline{z}_{i+1}^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)} \underline{z}_0^{(\ell)} \cdots \underline{z}_j^{(\ell)}$ for some $0 \leq j < i \leq n$.

(viii) Let $B = \underline{z}_0^{(\ell)}\, \underline{z}_1^{(\ell)} \cdots \underline{z}_{n-1}^{(\ell)}$ be a standard representative of a band of type 0 on level ℓ, and let $\omega \in \{0, 1, \dots, n-1\}$. We say B *allows a wrap-around on level* ℓ *at* ω if $\underline{z}_i^{(\ell)} = \underline{z}_{\omega-i}^{(\ell)}$ for all i, where the indices are taken modulo n.

Remark 4.2. (i) Let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$. Then S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level 1. There is a maximal positive integer ℓ such that S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ.

Conversely, if S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ relative to the $(\ell-1)$-tuples $(i_1, \dots, i_{\ell-1})$ and $(a_1, \dots, a_{\ell-1})$, then S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level j for all $1 \le j \le \ell$ relative to the $(j-1)$-tuples $(i_1, \ldots, i_{j-1})$ and $(a_1, \ldots, a_{j-1})$.

(ii) Let B be a standard representative of a band of type 0. Then B is a standard representative of a band of type 0 on level 1. There is a maximal positive integer ℓ such that B lies in the equivalence class under $\sim_{r,(0)}$ of a standard representative of a band of type 0 on level ℓ.

Conversely, if B is a standard representative of a band of type 0 on level ℓ relative to the $(\ell-1)$-tuples $(i_1, \ldots, i_{\ell-1})$ and $(a_1, \ldots, a_{\ell-1})$, then B is a standard representative of a band of type 0 on level j for all $1 \le j \le \ell$ relative to the $(j-1)$-tuples $(i_1, \ldots, i_{j-1})$ and $(a_1, \ldots, a_{j-1})$.

Proposition 4.3. *Let* $S = \beta\alpha^{-1}\underline{z}_0 \cdots \underline{z}_{n-1}\,\lambda$ *be the standard representative of a string of type* $\beta\alpha^{-1}\lambda$ *as in Definition 3.4. Then* S *has property* $(+)$ *from Lemma 3.6 if and only if there exist a positive integer* ℓ *and a choice of* $(\ell-1)$*-tuples* $(i_1, \ldots, i_{\ell-1})$ *and* $(a_1, \ldots, a_{\ell-1})$ *as in Definition 4.1(v) such that*

$$S = \underline{b}^{(\ell)}\,\underline{l}^{(\ell)} \quad \text{or} \quad S = \underline{b}^{(\ell)}\,(\underline{x}^{(\ell)})^{a_\ell}\,\underline{l}^{(\ell)} \quad \text{or} \quad S = \underline{b}^{(\ell)}\,(\underline{y}^{(\ell)})^{a_\ell}\,\underline{l}^{(\ell)} \tag{4.1}$$

for some positive integer a_ℓ.

Proof. Let ℓ be a positive integer, let $(i_1, \ldots, i_{\ell-1})$ and $(a_1, \ldots, a_{\ell-1})$ be $(\ell-1)$-tuples as in Definition 4.1(v), and let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ relative to these $(\ell-1)$-tuples. As stated in Remark 4.2(i), S is then also the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level j for all $1 \le j \le \ell$ relative to the $(j-1)$-tuples $(i_1, \ldots, i_{j-1})$ and $(a_1, \ldots, a_{j-1})$. Let $j \in \{1, \ldots, \ell\}$. Then $S = \underline{b}^{(j)}\,\underline{z}_0^{(j)} \cdots \underline{z}_{n_j-1}^{(j)}\,\underline{l}^{(j)}$, where $\underline{z}_i^{(j)} \in \{\underline{x}^{(j)}, \underline{y}^{(j)}\}$ for all $i \in \{0, \ldots, n_j - 1\}$. Let $\underline{c}^{(j)} = \underline{z}_0^{(j)}$ and let $\underline{d}^{(j)}$ be such that $\{\underline{c}^{(j)}, \underline{d}^{(j)}\} = \{\underline{x}^{(j)}, \underline{y}^{(j)}\}$. In particular, for $1 \le j < \ell$, $\underline{x}^{(j+1)} = (\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}$ and $\underline{y}^{(j+1)} = (\underline{c}^{(j)})^{a_j+1}\underline{d}^{(j)}$, and $\underline{b}^{(j+1)} = \underline{b}^{(j)}\,(\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}$ and $\underline{l}^{(j+1)} = (\underline{c}^{(j)})^{a_j}\underline{l}^{(j)}$.

For $1 \le j \le \ell$, consider the following property:

$(+)_j$ If there are subpatterns U, A_1, A_2 of S on level j such that

$$S = \underline{b}^{(j)}\,U\,\underline{c}^{(j)}A_2\,\underline{l}^{(j)} \ \text{ or } \ S = \underline{b}^{(j)}A_1\,\underline{c}^{(j)}\,U\,\underline{c}^{(j)}A_2\,\underline{l}^{(j)} \ \text{ or } \ S = \underline{b}^{(j)}A_1\,\underline{c}^{(j)}\,U\,\underline{l}^{(j)},$$

then for all subpatterns B_1, B_2 of S on level j, we have

$$S \ne \underline{b}^{(j)}\,U\,\underline{d}^{(j)}B_2\,\underline{l}^{(j)} \ \text{ and } \ S \ne \underline{b}^{(j)}B_1\,\underline{d}^{(j)}\,U\,\underline{d}^{(j)}B_2\,\underline{l}^{(j)} \ \text{ and } \ S \ne \underline{b}^{(j)}B_1\,\underline{d}^{(j)}\,U\,\underline{l}^{(j)}.$$

Claim 1. Let S be the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ as in the first paragraph of the proof. Then S has property $(+)$ from Lemma 3.6 if and only if S has property $(+)_\ell$.

Proof of Claim 1. Note that $(+)_1$ is the same as $(+)$. Moreover, S does not have property $(+)_j$ for some $1 \le j \le \ell$ if and only if there exist subpatterns U, A_1, A_2, B_1, B_2 of S on level j satisfying $(*)_j$ and $(**)_j$, where

$(*)_j$ $S = \underline{b}^{(j)}\,U\,\underline{c}^{(j)}A_2\,\underline{l}^{(j)}$ or $S = \underline{b}^{(j)}A_1\,\underline{c}^{(j)}\,U\,\underline{c}^{(j)}A_2\,\underline{l}^{(j)}$ or $S = \underline{b}^{(j)}A_1\,\underline{c}^{(j)}\,U\,\underline{l}^{(j)}$,

$(**)_j$ $S = \underline{b}^{(j)}\,U\,\underline{d}^{(j)}B_2\,\underline{l}^{(j)}$ or $S = \underline{b}^{(j)}B_1\,\underline{d}^{(j)}\,U\,\underline{d}^{(j)}B_2\,\underline{l}^{(j)}$ or $S = \underline{b}^{(j)}B_1\,\underline{d}^{(j)}\,U\,\underline{l}^{(j)}$.

Suppose that S does not have property $(+_j)$, i.e. S satisfies $(*)_j$ and $(**)_j$.

If $j < \ell$, then S is also the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level $(j+1)$. Therefore, it follows that the subpattern U in $(*)_j$ and $(**)_j$ has the form $U = (\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}\,\tilde{U}\,(\underline{c}^{(j)})^{a_j}$ for some subpattern $\tilde{U}$ of S on level $(j+1)$. Since $\underline{b}^{(j+1)} = \underline{b}^{(j)}\,(\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}$, $\underline{l}^{(j+1)} = (\underline{c}^{(j)})^{a_j}\underline{l}^{(j)}$, $\underline{x}^{(j+1)} = (\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}$ and $\underline{y}^{(j+1)} = (\underline{c}^{(j)})^{a_j+1}\underline{d}^{(j)}$, it follows that S satisfies $(*)_{j+1}$ and $(**)_{j+1}$ for appropriate subpatterns of S on level $(j+1)$.

If $j > 1$, then S is also the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level $(j-1)$. We have $\underline{b}^{(j)} = \underline{b}^{(j-1)}\,(\underline{c}^{(j-1)})^{a_{j-1}}\underline{d}^{(j-1)}$ and $\underline{l}^{(j)} = (\underline{c}^{(j-1)})^{a_{j-1}}\underline{l}^{(j-1)}$. Moreover, $\underline{c}^{(j)} = (\underline{c}^{(j-1)})^{\xi}\underline{d}^{(j-1)}$ and $\underline{d}^{(j)} = (\underline{c}^{(j-1)})^{\zeta}\underline{d}^{(j-1)}$, where $\{\xi, \zeta\} = \{a_{j-1}, a_{j-1}+1\}$. Using this to rewrite $(*)_j$ and $(**)_j$ on level $(j-1)$, it follows that S satisfies $(*)_{j-1}$ and $(**)_{j-1}$ for appropriate subpatterns of S on level $(j-1)$. Therefore, Claim 1 follows by induction.

Let now $S = \beta\alpha^{-1}\underline{z}_0\cdots\underline{z}_{n-1}\,\lambda$ be the standard representative of a string of type $\beta\alpha^{-1}\lambda$, i.e. S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level 1, and suppose S has property $(+)$ from Lemma 3.6. Using the notation from the first paragraph of the proof, it follows that $S = \underline{b}^{(1)}\,\underline{l}^{(1)}$, or $S = \underline{b}^{(1)}\,(\underline{c}^{(1)})^{a_1}\underline{l}^{(1)}$ for some positive integer a_1, or

$$S = \underline{b}^{(1)}\,(\underline{c}^{(1)})^{s_1}(\underline{d}^{(1)})^{t_1}\,\cdots\,(\underline{c}^{(1)})^{s_r}(\underline{d}^{(1)})^{t_r}(\underline{c}^{(1)})^{s_{r+1}}\,\underline{l}^{(1)}$$

for some positive integers r, $s_1,\ldots,s_{r+1}$ and $t_1,\ldots,t_r$ with $s_i = s_{r+2-i}$ and $t_j = t_{r+1-j}$ for all i,j by Remark 3.7. Since S has property $(+)$, it cannot contain both subpatterns $\underline{b}^{(1)}\,\underline{c}^{(1)}$ and $\underline{d}^{(1)}\,\underline{d}^{(1)}$. Therefore, $t_j = 1$ for all j. Because S cannot contain both subpatterns $\underline{b}^{(1)}\,(\underline{c}^{(1)})^{s_i}\underline{c}^{(1)}$ and $\underline{d}^{(1)}\,(\underline{c}^{(1)})^{s_i}\underline{d}^{(1)}$ (resp. S cannot contain both subpatterns $\underline{b}^{(1)}\,(\underline{c}^{(1)})^{s_1}\underline{d}^{(1)}$ and $\underline{c}^{(1)}\,(\underline{c}^{(1)})^{s_1}\underline{c}^{(1)}$), we obtain $s_1 \le s_i \le s_1+1$ for all i. Therefore,

$$S = \underline{b}^{(1)}\,(\underline{c}^{(1)})^{s_1}\,\underline{d}^{(1)}\,\cdots\,(\underline{c}^{(1)})^{s_r}\,\underline{d}^{(1)}\,(\underline{c}^{(1)})^{s_{r+1}}\,\underline{l}^{(1)} \tag{4.2}$$

for some positive integers r, $s_1,\ldots,s_{r+1}$ with $s_i \in \{s_1, s_1+1\}$ and $s_i = s_{r+2-i}$ for all i. Letting $a_1 = s_1$, we obtain that

$$S = \underline{b}^{(1)}\,(\underline{c}^{(1)})^{a_1}\,\underline{d}^{(1)}\,\underline{v}_0\cdots\underline{v}_{n_2-1}\,(\underline{c}^{(1)})^{a_1}\,\underline{l}^{(1)}$$

where $n_2 = r-1 \ge 0$ and $\underline{v}_i \in \{(\underline{c}^{(1)})^{a_1}\,\underline{d}^{(1)}, (\underline{c}^{(1)})^{a_1+1}\,\underline{d}^{(1)}\}$ for all $i \in \{0,\ldots,n_2-1\}$. Letting $i_1 = 0$ (resp. $i_1 = 1$) if $\underline{c}^{(1)} = \underline{x}$ (resp. $\underline{c}^{(1)} = \underline{y}$), we obtain

$$S = \underline{b}^{(2)}\,\underline{z}_0^{(2)}\cdots\underline{z}_{n_2-1}^{(2)}\,\underline{l}^{(2)} \tag{4.3}$$

where $n_2 \ge 0$ and $\underline{z}_i^{(2)} \in \{\underline{x}^{(2)}, \underline{y}^{(2)}\}$ for all $i \in \{0,\ldots,n_2-1\}$. If $n_2 \ge 1$, let $\underline{c}^{(2)} = \underline{z}_0^{(2)}$ and choose $\underline{d}^{(2)}$ such that $\{\underline{c}^{(2)}, \underline{d}^{(2)}\} = \{\underline{x}^{(2)}, \underline{y}^{(2)}\}$. Since S satisfies $(+)$, it also satisfies $(+)_2$ by Claim 1. Therefore, $S = \underline{b}^{(2)}\,\underline{l}^{(2)}$, or $S = \underline{b}^{(2)}\,(\underline{c}^{(2)})^{a_2}\underline{l}^{(2)}$ for some positive integer a_2, or

$$S = \underline{b}^{(2)}\,(\underline{c}^{(2)})^{u_1}(\underline{d}^{(2)})^{v_1}\,\cdots\,(\underline{c}^{(2)})^{u_w}(\underline{d}^{(2)})^{v_w}(\underline{c}^{(2)})^{u_{w+1}}\,\underline{l}^{(2)}$$

for some positive integers w, $u_1,\ldots,u_{w+1}$ and $v_1,\ldots,v_w$ with $u_i = u_{w+2-i}$ and $v_j = v_{w+1-j}$ for all i,j. Using induction, it follows that there exist a positive integer ℓ and a choice of $(\ell-1)$-tuples $(i_1,\ldots,i_{\ell-1})$ and $(a_1,\ldots,a_{\ell-1})$ as in Definition 4.1(v) such that S has one of the forms in (4.1).

Conversely, suppose that S is the standard representative of a string of type $\beta\alpha^{-1}\lambda$ on level ℓ having one of the forms in (4.1). Using the notation from the first paragraph of the proof, this means that $S = \underline{b}^{(\ell)}\,\underline{l}^{(\ell)}$, or $S = \underline{b}^{(\ell)}\,(\underline{c}^{(\ell)})^{a_\ell}\underline{l}^{(\ell)}$ for some positive integer a_ℓ. Hence S obviously has property $(+)_\ell$. By Claim 1, it follows that S has property $(+)$ from Lemma 3.6. This completes the proof of Proposition 4.3. □

Proposition 4.4. *Let $B = \underline{z}_0\,\underline{z}_1\,\cdots\,\underline{z}_{n-1}$ be a standard representative of a band of type 0 as in Definition 3.12. Then B has property $(++)$ from Lemma 3.14 if and only if there exist a positive integer ℓ and a choice of $(\ell-1)$-tuples $(i_1,\ldots,i_{\ell-1})$ and $(a_1,\ldots,a_{\ell-1})$ as in Definition 4.1(vi) such that*

$$B \sim_{r,(0)} \underline{x}^{(\ell)} \quad \text{or} \quad B \sim_{r,(0)} \underline{y}^{(\ell)}. \tag{4.4}$$

Moreover, if B has property $(++)$, then B allows exactly one wrap-around, as defined in Definition 3.12(iv).

Proof. Let ℓ be a positive integer, let $(i_1,\ldots,i_{\ell-1})$ and $(a_1,\ldots,a_{\ell-1})$ be $(\ell-1)$-tuples as in Definition 4.1(vi), and let B be a standard representative of a band of type 0 on level ℓ relative to these $(\ell-1)$-tuples. As stated in Remark 4.2(ii), B is then also a standard representative of a band of type 0 on level j for all $1 \le j \le \ell$ relative to the $(j-1)$-tuples $(i_1,\ldots,i_{j-1})$ and $(a_1,\ldots,a_{j-1})$. Let $j \in \{1,\ldots,\ell\}$. Then $B = \underline{z}_0^{(j)}\cdots\underline{z}_{n_j-1}^{(j)}$, where $\underline{z}_i^{(j)} \in \{\underline{x}^{(j)}, \underline{y}^{(j)}\}$ for all $i \in \{0,\ldots,n_j-1\}$. For $1 \le j < \ell$, define $\underline{c}^{(j)}, \underline{d}^{(j)} \in \{\underline{x}^{(j)}, \underline{y}^{(j)}\}$ such that $\underline{x}^{(j+1)} = (\underline{c}^{(j)})^{a_j}\underline{d}^{(j)}$ and $\underline{y}^{(j+1)} = (\underline{c}^{(j)})^{a_j+1}\underline{d}^{(j)}$.

For $1 \le j \le \ell$, consider the following property:

$(++)_j$ If there are rotated subpatterns U, V of B on level j such that $B \sim_{r,(0,j)} \underline{c}^{(j)}\, U\, \underline{c}^{(j)}\, V$, then for all rotated subpatterns W of B on level j, we have $B \not\sim_{r,(0,j)} \underline{d}^{(j)}\, U\, \underline{d}^{(j)}\, W$.

Claim 1. Let B be a standard representative of a band of type 0 on level ℓ as in the first paragraph of the proof. Then B has property $(++)$ from Lemma 3.14 if and only if B has property $(++)_\ell$.

Proof of Claim 1. Note that $(++)_1$ is the same as $(++)$. Moreover, B does not have property $(++)_j$ for some $1 \le j \le \ell$ if and only if there exist rotated subpatterns U, V, W of B on level j such that

$(\dagger\dagger)_j$ $B \sim_{r,(0,j)} \underline{c}^{(j)}\, U\, \underline{c}^{(j)}\, V$ and $B \sim_{r,(0,j)} \underline{d}^{(j)}\, U\, \underline{d}^{(j)}\, W$.

Suppose that B does not have property $(++)_j$, i.e. B satisfies $(\dagger\dagger)_j$.

If $j < \ell$, then B is also a standard representative of a band of type 0 on level $(j+1)$. Therefore, it follows that the rotated subpattern U in $(\dagger\dagger)_j$ has the form $U = (\underline{c}^{(j)})^{a_j}\, \underline{d}^{(j)}\, \tilde{U}\, (\underline{c}^{(j)})^{a_j}$ for some rotated subpattern $\tilde{U}$ of B on level $(j+1)$. Using that $\underline{x}^{(j+1)} = (\underline{c}^{(j)})^{a_j} \underline{d}^{(j)}$ and $\underline{y}^{(j+1)} = (\underline{c}^{(j)})^{a_j+1} \underline{d}^{(j)}$, it follows that B satisfies $(\dagger\dagger)_{j+1}$ for appropriate rotated subpatterns of B on level $(j+1)$.

If $j > 1$, then B is also a standard representative of a band of type 0 on level $(j-1)$. We have $\underline{c}^{(j)} = (\underline{c}^{(j-1)})^{\xi} \underline{d}^{(j-1)}$ and $\underline{d}^{(j)} = (\underline{c}^{(j-1)})^{\zeta} \underline{d}^{(j-1)}$, where $\{\xi, \zeta\} = \{a_{j-1}, a_{j-1}+1\}$. Using this to rewrite $(\dagger\dagger)_j$ on level $(j-1)$, it follows that B satisfies $(\dagger\dagger)_{j-1}$ for appropriate rotated subpatterns of B on level $(j-1)$. Therefore, Claim 1 follows by induction.

Let B be a standard representative of a band of type 0 on level ℓ having one of the forms in (4.4). Then B obviously has property $(++)_\ell$. By Claim 1, it follows that B has property $(++)$.

Conversely, suppose that $B = \underline{z}_0\, \underline{z}_1 \cdots \underline{z}_{n-1}$ is a standard representative of a band of type 0, i.e. B is a standard representative of a band of type 0 on level 1, and suppose B has property $(++)$. Then $B = \underline{x}^{(1)}$, or $B = \underline{y}^{(1)}$, or

$$B \sim_{r,(0)} (\underline{x}^{(1)})^{s_1} (\underline{y}^{(1)})^{t_1} \cdots (\underline{x}^{(1)})^{s_m} (\underline{y}^{(1)})^{t_m}$$

for some positive integers m, $s_1, \ldots, s_m$ and $t_1, \ldots, t_m$. Since B has property $(++)$, it cannot contain both rotated subpatterns $\underline{x}^{(1)}\, \underline{x}^{(1)}$ and $\underline{y}^{(1)}\, \underline{y}^{(1)}$. Therefore, $s_j = 1$ for all $1 \le j \le m$ or $t_j = 1$ for all $1 \le j \le m$. Let $(\underline{c}^{(1)}, \underline{d}^{(1)}) = (\underline{x}^{(1)}, \underline{y}^{(1)})$ and $e_j = s_j$ for $1 \le j \le m$ if $t_j = 1$ for all j, and let $(\underline{c}^{(1)}, \underline{d}^{(1)}) = (\underline{y}^{(1)}, \underline{x}^{(1)})$ and $e_j = t_j$ for $1 \le j \le m$ if $s_j = 1$ for all j. Then

$$B \sim_{r,(0)} (\underline{c}^{(1)})^{e_1}\, \underline{d}^{(1)} \cdots (\underline{c}^{(1)})^{e_m}\, \underline{d}^{(1)} \tag{4.5}$$

for some positive integers m, $e_1, \ldots, e_m$. Let a_1 be minimal among $e_1, \ldots, e_m$. Because B cannot contain both rotated subpatterns $\underline{c}^{(1)}\, (\underline{c}^{(1)})^{a_1} \underline{c}^{(1)}$ and $\underline{d}^{(1)}\, (\underline{c}^{(1)})^{a_1} \underline{d}^{(1)}$ for any $j \in \{1, \ldots, m\}$, it follows that $e_j \le a_1 + 1$ for all $1 \le j \le m$. Thus $e_j \in \{a_1, a_1+1\}$ for all $1 \le j \le m$, and we obtain that

$$B \sim_{r,(0)} \underline{v}_0\, \underline{v}_1 \cdots \underline{v}_{n_2-1}$$

where $n_2 = m \ge 1$ and $\underline{v}_i \in \{(\underline{c}^{(1)})^{a_1}\, \underline{d}^{(1)}, (\underline{c}^{(1)})^{a_1+1}\, \underline{d}^{(1)}\}$ for all $i \in \{0, \ldots, n_2 - 1\}$. Letting $i_1 = 0$ (resp. $i_1 = 1$) if $\underline{c}^{(1)} = \underline{x}$ (resp. $\underline{c}^{(1)} = \underline{y}$), we obtain

$$B \sim_{r,(0)} \underline{z}_0^{(2)}\, \underline{z}_1^{(2)} \cdots \underline{z}_{n_2-1}^{(2)} \tag{4.6}$$

where $n_2 \ge 1$ and $\underline{z}_i^{(2)} \in \{\underline{x}^{(2)}, \underline{y}^{(2)}\}$ for all $i \in \{0, \ldots, n_2 - 1\}$. Since B satisfies $(++)$, it follows that $\underline{z}_0^{(2)}\, \underline{z}_1^{(2)} \cdots \underline{z}_{n_2-1}^{(2)}$ satisfies $(++)_2$ by Claim 1. We obtain that $B \sim_{r,(0)} \underline{x}^{(2)}$, or $B \sim_{r,(0)} \underline{y}^{(2)}$, or

$$B \sim_{r,(0)} (\underline{x}^{(2)})^{u_1} (\underline{y}^{(2)})^{v_1} \cdots (\underline{x}^{(2)})^{u_w} (\underline{y}^{(2)})^{v_w}$$

for some positive integers w, $u_1, \ldots, u_w$ and $v_1, \ldots, v_w$. Using induction, it follows that there exist a positive integer ℓ and a choice of $(\ell-1)$-tuples $(i_1, \ldots, i_{\ell-1})$ and $(a_1, \ldots, a_{\ell-1})$ as in Definition 4.1(vi) such that B has one of the forms in (4.4).

It remains to show that if B is a standard representative of a band of type 0 satisfying condition $(++)$, then B allows exactly one wrap-around. By what we have just proved, we can assume that $\rho_u^{(0)}(B) = \underline{x}^{(\ell)}$ or $\rho_u^{(0)}(B) = \underline{y}^{(\ell)}$ for some $0 \le u \le n-1$. Hence $\rho_u^{(0)}(B)$ allows exactly one wrap-around on level ℓ. We can rewrite $\rho_u^{(0)}(B)$ as a standard representative of a band of type 0 on level j for all $1 \le j \le \ell$. Let $1 < j \le \ell$, and assume by induction that $\rho_u^{(0)}(B)$ allows exactly one wrap-around on level j. Using the notation from the first paragraph of the proof, it follows that

$$\rho_u^{(0)}(B) = \underline{z}_0^{(j)}\, \underline{z}_1^{(j)} \cdots \underline{z}_{n_j-1}^{(j)} \tag{4.7}$$

where $n_j \ge 1$ and $\underline{z}_i^{(j)} \in \{\underline{x}^{(j)}, \underline{y}^{(j)}\}$ for all $i \in \{0, \ldots, n_j - 1\}$. Moreover, $\underline{x}^{(j)} = (\underline{c}^{(j-1)})^{a_{j-1}}\underline{d}^{(j-1)}$ and $\underline{y}^{(j)} = (\underline{c}^{(j-1)})^{a_{j-1}+1}\underline{d}^{(j-1)}$. Therefore,

$$\begin{aligned} \rho_u^{(0)}(B) &= (\underline{c}^{(j-1)})^{u_0}\underline{d}^{(j-1)}\, (\underline{c}^{(j-1)})^{u_1}\underline{d}^{(j-1)} \cdots (\underline{c}^{(j-1)})^{u_{n_j-1}}\underline{d}^{(j-1)} \\ &= \underline{z}_0^{(j-1)}\, \underline{z}_1^{(j-1)} \cdots \underline{z}_{n_{j-1}-1}^{(j-1)} \end{aligned} \tag{4.8}$$

where $u_0, \ldots, u_{n_j-1} \in \{a_{j-1}, a_{j-1}+1\}$ and $n_{j-1} = (u_0+1)+\cdots+(u_{n_j-1}+1) = u_0+\cdots+u_{n_j-1}+n_j$. Assume that the unique wrap-around of $\rho_u^{(0)}(B)$ on level j is at w, so $\underline{z}_t^{(j)} = \underline{z}_{w-t}^{(j)}$ for all t, where the indices are taken modulo n_j. Let $\tilde{w} = (u_0+1)+\cdots+(u_{w-1}+1)+u_w-1$. Then $\rho_u^{(0)}(B)$ allows a wrap-around on level $(j-1)$ at $\tilde{w}$. On the other hand, if $\rho_u^{(0)}(B)$ allows a wrap-around on level $(j-1)$ at v, then (4.8) shows that v must be of the form $v = (u_0+1)+\cdots+(u_{\tilde{v}-1}+1)+u_{\tilde{v}}-1$ for some $\tilde{v} \in \{0, \ldots, n_j-1\}$. Hence this implies that $u_t = u_{\tilde{v}-t}$ for all t, and thus $\underline{z}_t^{(j)} = \underline{z}_{\tilde{v}-t}^{(j)}$ for all t, where the indices are taken modulo n_j. Therefore, the uniqueness of the wrap-around on level j leads to the uniqueness of the wrap-around on level $(j-1)$. Using induction, this completes the proof of Proposition 4.4. □

5. Appendix: The representation theory of Λ_0

As in §3, let k be an algebraically closed field of arbitrary characteristic and let Λ_0 be the basic k-algebra $\Lambda_0 = kQ/I$ where Q and I are as in Figure 1. In this appendix, we describe the representation theory of Λ_0. Since Λ_0 is a special biserial algebra, we can use the results from [7, 15]. It follows from [7] that all indecomposable non-projective Λ_0-modules are either string or band modules.

5.1. **String and band modules for** Λ_0**.** For each arrow $\alpha, \lambda, \xi, \delta, \rho, \beta$ in Q, we define a formal inverse $\alpha^{-1}, \lambda^{-1}, \xi^{-1}, \delta^{-1}, \rho^{-1}, \beta^{-1}$, respectively, with starting vertices $s(\alpha^{-1}) = 0 = s(\lambda^{-1})$, $s(\rho^{-1}) = 1 = s(\beta^{-1})$ and $s(\xi^{-1}) = 2 = s(\delta^{-1})$ and end vertices $e(\alpha^{-1}) = 0 = e(\beta^{-1})$, $e(\rho^{-1}) = 1 = e(\delta^{-1})$ and $e(\xi^{-1}) = 2 = e(\lambda^{-1})$. A word w is a sequence $w_1 \cdots w_n$, where w_i is either an arrow or a formal inverse such that $s(w_i) = e(w_{i+1})$ for $1 \le i \le n-1$. Define $s(w) = s(w_n)$, $e(w) = e(w_1)$ and $w^{-1} = w_n^{-1} \cdots w_1^{-1}$. There are also empty words 1_0, 1_1 and 1_2 of length 0 with $e(1_u) = u = s(1_u)$ and $(1_u)^{-1} = 1_u$ for $u \in \{0,1,2\}$. Denote the set of all words by $\mathcal{W}$, and the set of all non-empty words w with $e(w) = s(w)$ by $\mathcal{W}_r$. Let $J = \{\alpha^2, \rho^2, \xi^2, \alpha\lambda, \lambda\xi, \xi\delta, \delta\rho, \rho\beta, \beta\alpha, \lambda\delta\beta, \beta\lambda\delta, \delta\beta\lambda\}$.

Definition 5.1. Let $\sim_s$ be the equivalence relation on $\mathcal{W}$ with $w \sim_s w'$ if and only if $w = w'$ or $w^{-1} = w'$. Then *strings* are representatives $w \in \mathcal{W}$ of the equivalence classes under $\sim_s$ with the following property: Either $w = 1_u$ for $u \in \{0,1,2\}$, or $w = w_1 \cdots w_n$ where $w_i \ne w_{i+1}^{-1}$ for $1 \le i \le n-1$ and no subword of w or its formal inverse belongs to J.

Let $C = w_1 \cdots w_n$ be a string of length n. Then there exists an indecomposable Λ_0-module $M(C)$, called the *string module* corresponding to the string C, which can be described as follows. There is an ordered k-basis $\{z_0, z_1, \ldots, z_n\}$ of $M(C)$ such that the action of Λ_0 on $M(C)$ is given by the following representation $\varphi_C : \Lambda_0 \to \mathrm{Mat}(n+1, k)$. Let $v(i) = e(w_{i+1})$ for $0 \le i \le n-1$ and

$v(n) = s(w_n)$. Then for each vertex $u \in \{0,1,2\}$ and for each arrow $\zeta \in \{\alpha,\lambda,\xi,\delta,\rho,\beta\}$ in Q

$$\varphi_C(u)(z_i) = \left\{ \begin{array}{ccl} z_i & , & \text{if } v(i) = u \\ 0 & , & \text{else} \end{array} \right\} \text{ and } \varphi_C(\zeta)(z_i) = \left\{ \begin{array}{ccl} z_{i-1} & , & \text{if } w_i = \zeta \\ z_{i+1} & , & \text{if } w_{i+1} = \zeta^{-1} \\ 0 & , & \text{else} \end{array} \right\}.$$

We call φ_C the *canonical representation* and $\{z_0, z_1, \ldots, z_n\}$ a *canonical k-basis* for $M(C)$ relative to the representative C. Note that $M(C) \cong M(C^{-1})$.

The string modules for the empty words are isomorphic to the simple Λ_0-modules, namely $M(1_0) \cong S_0$, $M(1_1) \cong S_1$ and $M(1_2) \cong S_2$.

Definition 5.2. Let $w = w_1 \cdots w_n \in \mathcal{W}_r$. Then, for $0 \le i \le n-1$, the *i-th rotation* of w is defined to be the word $\rho_i(w) = w_{i+1} \cdots w_n w_1 \cdots w_i$. Let $\sim_r$ be the equivalence relation on $\mathcal{W}_r$ such that $w \sim_r w'$ if and only if $w = \rho_i(w')$ for some i or $w^{-1} = \rho_j(w')$ for some j. Then *bands* are representatives $w \in \mathcal{W}_r$ of the equivalence classes under $\sim_r$ with the following property: $w = w_1 \cdots w_n$, $n \ge 1$, with $w_i \ne w_{i+1}^{-1}$ and $w_n \ne w_1^{-1}$, such that w is not a power of a smaller word, and, for all positive integers m, no subword of w^m or its formal inverse belongs to J.

Let $B = w_1 \cdots w_n$ be a band of length n. Then for each integer $m > 0$ and each $\mu \in k^*$ there exists an indecomposable Λ_0-module $M(B,\mu,m)$ which is called the *band module* corresponding to the band B, μ and m, which can be described as follows. There is an ordered k-basis

$$\{z_{0,1}, z_{0,2}, \ldots, z_{0,m}, z_{1,1}, \ldots, z_{1,m}, \ldots, z_{n-1,1}, \ldots, z_{n-1,m}\}$$

of $M(B,\mu,m)$ such that the action of Λ_0 on $M(B,\mu,m)$ is given by the following representation $\varphi_{B,\mu,m} : \Lambda_0 \to \mathrm{Mat}(n{\cdot}m, k)$. Let $v(i) = e(w_{i+1})$ for $0 \le i \le n-1$. Then for each vertex $u \in \{0,1,2\}$ and for each arrow $\zeta \in \{\alpha,\lambda,\xi,\delta,\rho,\beta\}$ in Q, we have

$$\begin{aligned} \varphi_{B,\mu,m}(u)(z_{i,j}) &= \left\{ \begin{array}{ccl} z_{i,j} & , & \text{if } v(i) = u \\ 0 & , & \text{else} \end{array} \right\} \text{ and} \\ \varphi_{B,\mu,m}(\zeta)(z_{i,j}) &= \left\{ \begin{array}{ccl} \mu\, z_{0,j} + z_{0,j+1} & , & \text{if } w_i = \zeta \text{ and } i = 1 \\ z_{i-1,j} & , & \text{if } w_i = \zeta \text{ and } i \ne 1 \\ \mu^{-1} z_{1,j} + z_{1,j+1} & , & \text{if } w_{i+1} = \zeta^{-1} \text{ and } i = 0 \\ z_{i+1,j} & , & \text{if } w_{i+1} = \zeta^{-1} \text{ and } i \ne 0 \\ 0 & , & \text{else} \end{array} \right\} \end{aligned}$$

for all $0 \le i \le n-1$, $1 \le j \le m$, where $z_{0,m+1} = 0 = z_{1,m+1}$ and $z_{n,j} = z_{0,j}$ for all j. We call $\varphi_{B,\mu,m}$ the *canonical representation* and $\{z_{0,1}, z_{0,2}, \ldots, z_{0,m}, z_{1,1}, \ldots, z_{1,m}, \ldots, z_{n-1,1}, \ldots, z_{n-1,m}\}$ a *canonical k-basis* for $M(B,\mu,m)$ relative to the representative B. Note that we have for all $0 \le i,j \le n-1$,

$$M(B,\mu,m) \cong M(\rho_i(B),\mu,m) \cong M(\rho_j(B)^{-1},\mu^{-1},m).$$

5.2. **The stable Auslander-Reiten quiver of Λ_0.** Each component of the stable Auslander-Reiten quiver of Λ_0 consists either entirely of string modules or entirely of band modules. The band modules all lie in 1-tubes. The components consisting of string modules are two 3-tubes and infinitely many non-periodic components of type $\mathbb{Z}A_\infty^\infty$.

The irreducible morphisms between string modules can be described using hooks and cohooks. For our algebra Λ_0, these are defined as follows. Let $\mathcal{M}$ be the set of maximal directed strings, i.e. $\mathcal{M} = \{\alpha, \rho, \xi, \lambda\delta, \delta\beta, \beta\lambda\}$.

Definition 5.3. Let S be a string. We say that S *starts on a peak* (resp. *starts in a deep*) if $S = S'C$ (resp. $S = S'C^{-1}$) for some string C in $\mathcal{M}$. Dually, we say that S *ends on a peak* (resp. *ends in a deep*) if $S = D^{-1}S''$ (resp. $S = DS''$) for some string D in $\mathcal{M}$.

If S does not start on a peak (resp. does not start in a deep), there is a unique arrow ζ and a unique $M \in \mathcal{M}$ such that $S_h = S\zeta M^{-1}$ (resp. $S_c = S\zeta^{-1}M$) is a string. We say S_h (resp. S_c) is obtained from S by adding a *hook* (resp. a *cohook*) on the right side.

Dually, if S does not end on a peak (resp. does not end in a deep), there is a unique arrow ξ and a unique $N \in \mathcal{M}$ such that ${}_hS = N\xi^{-1}S$ (resp. ${}_cS = N^{-1}\xi S$) is a string. We say ${}_hS$ (resp. ${}_cS$) is obtained from S by adding a *hook* (resp. a *cohook*) on the left side.

All irreducible morphisms between string modules are either canonical injections $M(S) \to M(S_h)$, $M(S) \to M({}_hS)$, or canonical projections $M(S_c) \to M(S)$, $M({}_cS) \to M(S)$.

In particular, since none of the projective Λ_0-modules is uniserial, we get the following result. Suppose $M(S)$ is a string module of minimal length such that $M(S)$ belongs to a component of the stable Auslander-Reiten quiver of Λ_0 of type $\mathbb{Z}A_\infty^\infty$. Then near $M(S)$ the stable Auslander-Reiten component looks as in Figure 2.

FIGURE 2. The stable Auslander-Reiten component near $M(S)$.

$$\begin{tikzcd}
 & & & & & & \\
 & M({}_{cc}S) \arrow[dr] \arrow[ul, dotted, no head] \arrow[ur, dotted, no head] \arrow[dl, dotted, no head] & & M({}_cS_h) \arrow[dr] \arrow[ul, dotted, no head] \arrow[ur, dotted, no head] & & M(S_{hh}) \arrow[ul, dotted, no head] \arrow[ur, dotted, no head] \arrow[dr, dotted, no head] & \\
 & & M({}_cS) \arrow[ur] \arrow[dr] & & M(S_h) \arrow[ur] \arrow[dr] & & \\
 & M({}_cS_c) \arrow[ur] \arrow[dr] \arrow[ul, dotted, no head] \arrow[dl, dotted, no head] & & M(S) \arrow[ur] \arrow[dr] & & M({}_hS_h) \arrow[ur, dotted, no head] \arrow[dr, dotted, no head] & \\
 & & M(S_c) \arrow[ur] \arrow[dr] & & M({}_hS) \arrow[ur] \arrow[dr] & & \\
 & M(S_{cc}) \arrow[ur] \arrow[ul, dotted, no head] \arrow[dl, dotted, no head] \arrow[dr, dotted, no head] & & M({}_hS_c) \arrow[ur] \arrow[dl, dotted, no head] \arrow[dr, dotted, no head] & & M({}_{hh}S) \arrow[dl, dotted, no head] \arrow[dr, dotted, no head] \arrow[ur, dotted, no head] & \\
 & & & & & &
\end{tikzcd}$$

5.3. **Homomorphisms between string and band modules for Λ_0.** In [15], all homomorphisms between string and band modules have been determined. The following remark describes the homomorphisms between string modules using the canonical bases defined in Definition 5.1.

Remark 5.4. Let $M(S)$ (resp. $M(T)$) be a string module for Λ_0 with a canonical k-basis $\{x_u\}_{u=0}^m$ (resp. $\{y_v\}_{v=0}^n$) relative to the representative S (resp. T). Suppose C is a string such that

(i) $S \sim_s S'CS''$ with (S' of length 0 or $S' = \hat{S}'\xi_1$) and (S'' of length 0 or $S'' = \xi_2^{-1}\hat{S}''$), where $S', \hat{S}', S'', \hat{S}''$ are strings and ξ_1, ξ_2 are arrows in Q; and

(ii) $T \sim_s T'CT''$ with (T' of length 0 or $T' = \hat{T}'\zeta_1^{-1}$) and ($T''$ of length 0 or $T'' = \zeta_2\hat{T}''$), where $T', \hat{T}', T'', \hat{T}''$ are strings and ζ_1, ζ_2 are arrows in Q.

Then by [15] there exists a non-zero Λ_0-module homomorphism $\sigma_C : M(S) \to M(T)$ which factors through $M(C)$ and which sends each element of $\{x_u\}_{u=0}^m$ either to zero or to an element of $\{y_v\}_{v=0}^n$, according to the relative position of C in S and T, respectively. If e.g. $S = s_1s_2\cdots s_m$, $T = t_1t_2\cdots t_n$, and $C = s_{i+1}s_{i+2}\cdots s_{i+\ell} = t_{j+\ell}^{-1}t_{j+\ell-1}^{-1}\cdots t_{j+1}^{-1}$, then

$$\sigma_C(x_{i+t}) = y_{j+\ell-t} \text{ for } 0 \le t \le \ell, \text{ and } \sigma_C(x_u) = 0 \text{ for all other } u.$$

We call σ_C a *canonical homomorphism* from $M(S)$ to $M(T)$. Note that there may be several choices for S', S'' (resp. T', T'') in (i) (resp. (ii)). In other words, there may be several k-linearly independent canonical homomorphisms factoring through $M(C)$. By [15], every Λ_0-module homomorphism $\sigma : M(S) \to M(T)$ can be written as a unique k-linear combination of canonical homomorphisms which factor through string modules corresponding to strings C satisfying (i) and (ii).

It follows from [15] that if B is a band, $\mu \in k^*$ and $n \geq 2$ is an integer, then $\underline{\mathrm{End}}_{\Lambda_0}(M(B,\mu,n))$ has k-dimension at least 2. The following remark describes the homomorphisms between band modules of the form $M(B,\mu,1)$ using the canonical bases defined in Definition 5.2.

Remark 5.5. Let $B, \tilde{B}$ be bands and let $\mu, \tilde{\mu} \in k^*$. Let $M_{B,\mu} = M(B,\mu,1)$ (resp. $M_{\tilde{B},\tilde{\mu}} = M(\tilde{B},\tilde{\mu},1)$) with a canonical k-basis $\{z_{u,1}\}_{0\leq u\leq m-1}$ (resp. $\{\tilde{z}_{v,1}\}_{0\leq v\leq n-1}$) relative to the representative B (resp. $\tilde{B}$). Suppose S is a string such that

(i) $B \sim_r ST_1$ with $T_1 = \xi_1^{-1}T_1'\xi_2$, where T_1, T_1' are strings and ξ_1, ξ_2 are arrows in Q; and
(ii) $\tilde{B} \sim_r ST_2$ with $T_2 = \zeta_1 T_2' \zeta_2^{-1}$, where T_2, T_2' are strings and ζ_1, ζ_2 are arrows in Q.

Then by [15] there exists a non-zero Λ_0-module homomorphism $\tau_S : M_{B,\mu} \to M_{\tilde{B},\tilde{\mu}}$ which factors through $M(S)$ and which sends each element of $\{z_{u,1}\}_{0\leq u\leq m-1}$ either to zero or to an element of $\{\tilde{z}_{v,1}\}_{0\leq v\leq n-1}$, according to the relative position of S in B and $\tilde{B}$, respectively. Suppose e.g. $B = w_1\cdots w_m$, $\tilde{B} = \tilde{w}_1\cdots\tilde{w}_n$ and $S = w_{i+1}w_{i+2}\cdots w_{i+\ell} = \tilde{w}_{j+\ell}^{-1}\tilde{w}_{j+\ell-1}^{-1}\cdots\tilde{w}_{j+1}^{-1}$, where $w_i = \xi_2$, $w_{i+\ell+1} = \xi_1^{-1}$, $\tilde{w}_j^{-1} = \zeta_1$, $\tilde{w}_{j+\ell+1}^{-1} = \zeta_2^{-1}$. Then

$$\tau_S(z_{i+t,1}) = \tilde{z}_{j+\ell-t,1} \text{ for } 0 \leq t \leq \ell, \text{ and } \tau_S(z_{u,1}) = 0 \text{ for all other } u.$$

We call τ_C a *canonical homomorphism* from $M_{B,\mu}$ to $M_{\tilde{B},\tilde{\mu}}$ *of string type* S. Note that there may be several choices for T_1 (resp. T_2) in (i) (resp. (ii)). In other words, there may be several k-linearly independent canonical homomorphisms of string type S. By [15], if $M_{B,\mu}$ and $M_{\tilde{B},\tilde{\mu}}$ are not isomorphic, then every Λ_0-module homomorphism $\tau : M_{B,\mu} \to M_{\tilde{B},\tilde{\mu}}$ can be written as a unique k-linear combination of canonical homomorphisms of string type S for suitable choices of strings S satisfying (i) and (ii). If $B = \tilde{B}$ and $\mu = \tilde{\mu}$, then every Λ_0-module endomorphism of $M_{B,\mu}$ can be written as a unique k-linear combination of the identity homomorphism and of canonical endomorphisms of string type S for suitable choices of strings S satisfying (i) and (ii).

REFERENCES

[1] F. M. Bleher, Universal deformation rings and Klein four defect groups. Trans. Amer. Math. Soc. 354 (2002), 3893–3906.

[2] F. M. Bleher, Universal deformation rings and dihedral defect groups. Trans. Amer. Math. Soc. 361 (2009), 3661–3705.

[3] F. M. Bleher and T. Chinburg, Universal deformation rings and cyclic blocks. Math. Ann. 318 (2000), 805–836.

[4] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. C. R. Math. Acad. Sci. Paris 342 (2006), 229–232.

[5] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. Math. Ann. 337 (2007), 739–767.

[6] G. Böckle, Presentations of universal deformation rings. In: L-functions and Galois representations, 24–58, London Math. Soc. Lecture Note Ser., 320, Cambridge Univ. Press, Cambridge, 2007.

[7] M. C. R. Butler and C. M. Ringel, Auslander-Reiten sequences with few middle terms and applications to string algebras. Comm. Algebra 15 (1987), 145–179.

[8] T. Chinburg, Can deformation rings of group representations not be local complete intersections? In: Problems from the Workshop on Automorphisms of Curves. Edited by Gunther Cornelissen and Frans Oort, with contributions by I. Bouw, T. Chinburg, Cornelissen, C. Gasbarri, D. Glass, C. Lehr, M. Matignon, Oort, R. Pries and S. Wewers. Rend. Sem. Mat. Univ. Padova 113 (2005), 129–177.

[9] G. Cornell, J. H. Silverman and G. Stevens (eds.), Modular Forms and Fermat's Last Theorem (Boston, 1995). Springer-Verlag, Berlin-Heidelberg-New York, 1997.

[10] C. W. Curtis and I. Reiner, Methods of representation theory. Vol. I. With applications to finite groups and orders. John Wiley and Sons, Inc., New York, 1981.

[11] B. de Smit and H. W. Lenstra, Explicit construction of universal deformation rings. In: Modular Forms and Fermat's Last Theorem (Boston, MA, 1995), Springer-Verlag, Berlin- Heidelberg-New York, 1997, pp. 313–326.

[12] K. Erdmann, Blocks of Tame Representation Type and Related Algebras, Lecture Notes in Mathematics, vol. 1428, Springer-Verlag, Berlin-Heidelberg-New York, 1990.

[13] R. Ile, Change of rings in deformation theory of modules. Trans. Amer. Math. Soc. 356 (2004), 4873–4896.

[14] B. Keller, Hochschild cohomology and derived Picard groups. J. Pure Appl. Algebra Algebra 190 (2004), 177–196.

[15] H. Krause, Maps between tree and band modules. J. Algebra 137 (1991), 186–194.

[16] O. A. Laudal, Noncommutative deformations of modules. The Roos Festschrift volume, 2. Homology Homotopy Appl. 4 (2002), 357–396.

[17] B. Mazur, Deformation theory of Galois representations. In: Modular Forms and Fermat's Last Theorem (Boston, MA, 1995), Springer Verlag, Berlin-Heidelberg-New York, 1997, pp. 243–311.
[18] M. Schlessinger, Functors of Artin Rings. Trans. of the Amer. Math. Soc. 130, 1968, 208–222.
[19] J. A. Vélez-Marulanda, Universal deformation rings of modules over self-injective algebras, Dissertation, University of Iowa, 2010.
[20] D. Yau, Deformation theory of modules, Comm. Algebra 33 (2005), 2351–2359.

F.B.: Department of Mathematics, University of Iowa, 14 MacLean Hall, Iowa City, IA 52242-1419, U.S.A.
E-mail address: `frauke-bleher@uiowa.edu`

J.V.: Department of Mathematics & Computer Science, Valdosta State University, 2072 Nevins Hall, Valdosta, GA 31698-0040, U.S.A.
E-mail address: `javelezmarulanda@valdosta.edu`